



DEVELOPMENT AND VALIDATION OF RP-HPLC METHOD FOR ESTIMATION OF TRIMIPRAMINE MALEATE IN PHARMACEUTICAL DOSAGE FORM

Mr. Pravin Rajendra Maharnur*¹, Prof. S. A. Waghmare², Dr. Hemant Kambale³

¹Research Scholar, M. Pharm, Loknete shri Dada Patil Pharate college of Pharmacy, Mandavgan.

²Assistant Professor, Loknete shri Dada Patil Pharate college of Pharmacy, Mandavgan.

³Principal, Loknete shri Dada Patil Pharate college of Pharmacy, Mandavgan.

ABSTRACT

A simple, precise, rapid, accurate and economic reverse phase high performance liquid chromatographic method have been developed for the estimation of Trimipramine maleate in various pharmaceutical dosage form like tablet, capsule, gel, etc.

The selection of mobile phase as well as stationary phase is finalized after assessing the solubility of Trimipramine maleate in different organic solvent as well on the basis of literature survey. The Trimipramine maleate having anti-depressant activity, it is odourless, white crystalline powder. Also soluble in different organic solvent and sparingly soluble in aquatic buffer.

KEYWORDS: Trimipramine maleate, RP-HPLC, Validation,

INTRODUCTION

HPLC is an analytical process utilizing special instruments designed to separate, quantify and analyse components of chemical mixture. RP-HPLC is the choice for the majority of samples. It consists of a non-polar stationary phase and an aqueous, moderately polar mobile phase. One common stationary phase is silica which has been treated with alkyl dimethyl silylchloride (RMe₂SiCl), where R is a straight chain alkyl group such as octadecyl (C₁₈H₃₇) or octyl (C₈H₁₇). The retention time is therefore longer for molecules which are more non-polar in nature, allowing polar molecules to elute more readily.

By using the reverse phase HPLC method for the estimation of Trimipramine maleate in different pharmaceutical dosage form like Tablet, Capsule, etc. the new precise, economical, simple, rapid & accurate method are developed and validate. For the estimation chromatographic analysis method used. As per the study of different literature survey selection of column, Selection of stationary as well as mobile phase is good for accurate result. In the estimation of Trimipramine maleate the analysis of single dose formulation of Trimipramine maleate tablet can also be successfully performed by the RP-HPLC method. No interference of additives, matrix etc. is encountered in these methods.



Figure 1.1: Typical HPLC Waters System

DRUG PROFILE

Trimipramine Maleate Structure




Fig. 6.1 Trimipramine Maleate.
General profile of Trimipramine maleate

Category	Anti-depressant agent.
Chemical Name	3-(5,6-dihydrobenzo[b][1]benzazepin-11-yl)-N,N,2-trimethylpropan-1-amine
Molecular Formula	C ₂₀ H ₂₆ N ₂
Molecular Weight	294.434 g/mol
Odour	Odourless
Description	White crystalline powder.
Solubility	soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), it is sparingly soluble in aqueous buffers.
Melting point	45°C
Odour	Odourless
PKa	9.24 (Strongest Acidic)
LOG P-	4.2

Reported Analytical Method

After the study of different literature survey on drug Trimipramine maleate following analytical methods are found to be reported and research gap are also found.

Sr. No.	Name of Drug	Mobile Phase composition	Discussion	Ref
1	Trimipramine maleate	Methanol: acetonitrile (20:80, v/v)	The retention time of Trimipramine maleate was 10.4 min in plasma and 10.9 min in urine.	1
2	Trimipramine maleate	ammonium formate: Methanol (25:75V/V)	Retention time of Trimipramine maleate was 1.67Min for R configuration & 1.48 Min for S Configuration.	2
3	Trimipramine maleate	Methanol: water (60:40 % v/v)	The retention time of Trimipramine maleate was 9.1 min	3
4	Trimipramine maleate	sodium hydrogen phosphate solution -acetonitrile (60:40, v/v)	The retention times of Trimipramine maleate were 4.3 and 5.2 min. respectively	4
5	Trimipramine maleate	hexane/isoamyl alcohol 98:2 (v/v)	The retention time of Trimipramine maleate was 9.1 min.	5
6	Trimipramine maleate and Opipramol	ammonium formate with 0.1% formic acid:methanol 25:75(v/v)	The retention time of Trimipramine and Opipramol were found to be 1.67 min and 1.48 min respectively.	6



CONCLUSION

- The develop new, simple, sensitive, accurate, and economical analytical method for the determination of assay of Trimipramine maleate in capsule dosage form by RP-HPLC.
- Validate the proposed method in accordance with USP and ICH guidelines for the intended analytical application i.e., to apply the proposed method for analysis of the drug in its dosage form.
- The method shows good reproducibility; moreover, the RP-HPLC method is accurate, precise, specific, reproducible and sensitive.
- The analysis of single dose formulation of Trimipramine maleate tablet can also be successfully performed by the RP-HPLC method. No interference of additives, matrix etc. is encountered in these methods.
- Further studies on other pharmaceutical formulations would throw more light on these studies. Suitability of these methods on biological samples needy also studies.

REFERENCES

1. Meyer veronica r.; *practical high performance liquid chromatography*, john wiley and sons, london, 2nd edition, 1993, 26, 27, 40, 222, 246, 258.
2. Chatwal g. R., anand s. K.; *instrumental method of chemical analysis*; himalaya publishing house, 11th edition, 2005, 2.634-2.638
3. Raymond p. W. Scott; *liquid chromatography for the analyst*, chromatographic science series; marcel dekker, inc., 1991, 1-30.
4. Andrea westen; *hplc and ce – principles and practice*; academic press 1997, 1- 21.
5. Snyder l.r; *high-performance liquid chromatography: advances and perspectives*; c. Horvath, ed., academic press, san diego, ca; 1983, 3, 157.
6. Snyder l. R., m. A. Stadalius.; *high-performance liquid chromatography: advances and persepectives*; c. Horvath, ed., academic press, san diego, ca; 1986, 4, 294-295.
7. Ghulam a. Shabir.; *hplc method development and validation for pharmaceutical analysis*; pharmaceutical technology europe, march 2004.
8. Paul c. Sadek.; *troubleshooting hplc systems*; john wiley and sons, new york, 2000
9. [Http://www.waters.com/](http://www.waters.com/)
10. Sethi p.d.; *hplc-quantitative analysis of pharmaceutical formulation*; cbs publisher and distributors new delhi; 2001, 11.
11. Ich q2b harmonized tripartite guideline, *text on validation of analytical procedures*, ifpma, in proceeding of international conference on harmonization, geneva, march 1996.
12. D.a.skoog, f.j. Holler, a. Timothy, n.w.nieman, *principle of instrumental analysis*, saunders college publications, london, 1998.
13. Asean, *guideline on stability of drug product*, philippines, 2005.
14. S.w.baertschi, p.j. Jansen, *pharmaceutical stress testing*, by taylor&francis group, llc, 2005.
15. S. Klick, *stress testing of drug substances and drug products*, pharm. Technol. 2005.
16. Ich, *stability testing: photostability testing of new drug substances and products q1b*, international conference on harmonisation, ifpma, geneva, 1996.
17. [Www.drugsbank.com](http://www.drugsbank.com).
18. Dyade g.k., sharmaa.k.; *indian drugs*, 2001, 38(2): 75-78.
19. Sethi p.d.; *qualitativie analysis of drugs in pharmaceutical formulations*, 3rd edition, 1997, 182-184.
20. Swarbrick james.,boylanjames.c.; *encyclopedia of pharmaceutical technology*, volume i, marcel dekkerinc., new york, 1998, 217 - 224.
21. Lindsay sandy.; *hplc by open learning*; john wiley and sons, london, 1991, 30- 45.
22. Lough w.j., waineri.w.w.; *hplc fundamental principles and practices*, blackie academic and professional, 1991, 52-67.
23. G. D christian; in: *analytical chemistry*, 4th edition, john wiley and sons, United Kingdom, 1986, 1-6.