



COMPARISON OF HAEMATOLOGICAL AND BIOCHEMICAL PARAMETERS IN SHEEPS AND MICE BALB\C STRAIN INFECTED WITH HYDATID CYST

Muslim Abbas Allu

Department of Biology, University of Zakho, Ministry of Higher Education and Scientific Research, Kurdistan Region, Iraq.

ABSTRACT

The paper reviews Human hydatid cyst or cystic echinococcosis is a life-threatening zoonotic disease that occurs in most countries worldwide and is recognized as a major public health problem. Following, ingestion of *Echinococcus granulosus* eggs, hydatid cysts which is the larval stage of the worm are formed mostly in liver and lungs, and occasionally in other organs of human, the infection with hydatid disease effect on the liver and lung function after examination of some biochemical parameters that related to liver and lung function,. Aspartate Total Protein, Total Bilirubin, Aminotransferase (AST), Lactate Dehydrogenase (LDH) were decrease while aspartate Blood Urea Nitrogen (BUN), Creatinine, Albumin has increase, Values of hemoglobin and red blood cells and packed cell volume in sheep infested by *E. granulosus*, appear low compared with reference values. While hyper eosinophilia, lymphocyte and monocyte was high, and this typical of parasitic diseases. In mice Balb \c strain and sheep found that the number of white blood cell has increase (eosinophilia, lymphocyte and monocyte) and decrease of neutrophils after counting the differential WBC of positive group and increase of MCV, MCH, MCHC.

KEYWORDS: hydatid cyst, haematological parameters, biochemical indicators, sheeps, mice balb \c.

INTRODUCTION

Hydatid disease is a wide spread in all countries of the world, and for large numbers of dogs and adults' worms (1). It is one of the chronic diseases in humans to continue having a cyst for long time (2) Cystic echinococcosis is a zoonotic disease caused by the larvae of the *Echinococcus granulosus* (*E. granulosus*) infecting the intermediate hosts, and the disease represents a significant threat for human and animal health all around the globe (3,4,5,6). The disease infected by *E. granulosus* most often spreads to the liver (50%-70%) and less frequently the lung and in other parts of the body. (7)

EPIDEMIOLOGY

The hydatid cyst remains a significant public health hazard in endemic areas such as Iraq, Turkey, the Middle East, South America, New Zealand, Africa, China, northern Kenya, Australia, and other sheep-raising areas (8). As an endemic disease, it causes social and economic losses for countries. WHO reports stated that approximately 100,000 people in the world are infected with this disease every year (9) which is common in rural populations of underdeveloped countries because of their close association with domestic and wild animals (10). The incidence of infestation by *E. granulosus* in endemic areas ranges from 1 - 220 cases per 100,000 inhabitants. Diagnosis of hydatid disease is based on the epidemiological background of patients, clinical signs and symptoms, and radiographic and related imaging studies. Serodiagnostic assays can be particularly helpful and detection of antibody to specific echinococcal antigens has the highest degree of specificity.

PATHOGENESIS

Hydatid cysts are generally asymptomatic until expanding cysts gradually effect in the liver and elicit pressure symptoms. Cysts

may grow for a period of 5-20 years, and may be discovered incidentally on a routine ultrasound or CT examination. Expanding cysts present with abdominal pain or palpable mass in the right upper abdomen. (11) HCs may compress the bile ducts and result in obstructive jaundice and cause fever, pruritus, eosinophilia, dissemination, or fatal anaphylaxis after a cyst ruptures into the peritoneum or biliary tract. Infection of the cyst can facilitate the development of liver abscesses and mechanic local complications, such as mass effect on bile ducts and vessels that can induce cholestasis, portal hypertension, and Budd-Chiari syndrome. (12-16) As a result, the diagnosis of HCs should be as fast as possible because of the relevant complications that may arise with disease progression.

HOSTS AND TRANSMISION

A special feature of the liver, as a body, is the fact that its pathologies, in most cases, are not manifested clinically, and the damage that comes from their health and animal production, is great (17). Touching the body will affect systemic change and ultimately causes a decrease in production. The disease is known as many labels, including Hydatidosis, echinococcosis or cystic echinococcosis (18-21) Carnivorous definitive host for the parasite and include the canine (Dogs, hyenas, wolves, vany and some For other fierce animals) while animals are Herbivorous such as (sheep, cows, camels, buffalo, horses and others) Intermediate Host for parasite (22-24) The disease affects parts that exist in human and the intermediate host, especially liver and lung, followed by the rest of the parts, such as spleen, brain and other muscles except for hair and nails (25-28). The most important source of infection for humans and animals that are the intermediate hosts are the infected dogs. The vegetables, fruits, and drinking waters contaminated with the dog feces form the source of infection (29). The parasite is



reported to be capable of settling into almost every organ, including lungs, kidneys, spleen, brain, and heart (30).

DIAGNOSIS

Imaging studies, combined with immunodiagnostic techniques, often help to make a diagnosis (31,32). Ultrasonography (USG) is the initial imaging modality of choice because it is easy to perform, widely available, and inexpensive and can help define the number, site, size, and vitality of cysts (33). Antibody assays can add weight to the presumptive imaging diagnosis. However, a negative serologic test rarely rules out echinococcosis (34). Computed tomography (CT) scan and magnetic resonance imaging can help diagnose deep-seated lesions and determine the extent and condition of the avascular fluid-filled cysts (31,35).

All cases have hydatidosis disease according to identification of the parasite's structures by imaging techniques, including ultrasound, x-ray diagnosis and accentuated surgery operation of each patient. Cysts also contain 'daughter cysts' which, if released, may diffuse to other areas of the body (36). Many substances have recently been used to activate and modify the host's immune system in order to control the growth and development of the cyst (37). The clinical manifestation of Echinococcosis infection depends on the site and size of the cyst. The infection may remain asymptomatic in earlier stages when the cyst is small (38). Eventually, as the disease progresses, HCL can present with epigastric or right upper quadrant abdominal pain, nausea, vomiting, and hepatomegaly (39-40).

BLOOD PARAMETERS

In sheep and mice Balb/c strain injected by hydatid cyst protoscolexes. it was found that the numbers of WBC, lymphocytes, eosinophils and monocyte have increased while the number of neutrophils decreased in the positive group compared with the control group, these results agreed with the study of (45,46), who found an increase in the number of leukocytes, lymphocytes and eosinophils and a decrease in the number of neutrophils. They, however, disagree with (42) who found an increase in the number of neutrophils. Increases in numbers of WBC, lymphocytes and eosinophils were observed in the present study which may be considered as a defense mechanism against the inflammatory processes in the body especially in the liver, spleen and kidneys where the inflammation stimulates the bone marrow to produce a large number of WBC. The increase of the eosinophil count could be attributed to the long period of the disease (47) explained that eosinophilia was produced due to the ability of parasites to infect the tissue and this agreed with (40). PCV, HB, MCV, MCH and MCHC had decrease of positive group comparison with control groups in sheep and mice Balb/c strain injected by hydatid cyst protoscolexes. Our values compared to the reference, indicate for significant anemia of sheep. HB, PCV level was significantly lower than the reference value by (48,49,50). Elevated levels of eosinophils, neutrophils and lymphocytes particularly, and the early period of the infection encountered in this study is in conformity with the findings of (51) that elevated eosinophils, neutrophils and lymphocytes participate in the defense against *F. hepatica*. This significant

increase in the number of eosinophils, hipereosinophilia, shows high parasitic burden of these animals since the increase in the number of these cells in peripheral blood is characteristic of parasitic infestation and is cited by some authors (52). Significant changes reportedly occur in blood parameters of animals exposed to the parasitic invasion (53,54,55). Parasitic infections seem to cause increased hepatic enzyme activity and decreased trace element levels by causing liver damage (56-58). The determination of blood parameters is further important as the rate of success in the treatment increases with early diagnosis of the disease (59).

BIOCHEMICAL ANALYSIS

HCs also plays a definitive role in the metabolism, physiology and immunology of cystic echinococcosis (60), radiologic appearance of HCs depends on the stage of maturity and ranges from completely liquid type to completely solid type (61). Some researchers report that investigating the hepatic enzyme activities and total bilirubin, protein, and albumin levels are helpful in the evaluation of the liver functions (62), minerals, on the other hand, have an important role in maintaining normal physiological functions and protecting organisms against disease. substances are found in the structure of cofactors that These are necessary for enzymatic activity (63).

Biochemical profile in sheep infested naturally by CE and comparison these value with the biochemical value in mice Balb/c strain injected by hydatid cyst protoscolexes, and investigate the value of biochemical profile for infected group and comparison with control group. after examination of some biochemical parameters that related to liver and lung function, Aspartate Total Protein, Total Bilirubin, Aminotransferase (AST), and Lactate Dehydrogenase (LDH) had decrease while aspartate Blood Urea Nitrogen (BUN), Albumin and Creatinine, has increase, comparison with control group in sheep, While in mice aspartate Total Protein, Total Bilirubin, Aminotransferase (AST), and Lactate Dehydrogenase (LDH) had decrease while aspartate Blood Urea Nitrogen (BUN), Creatinine and Albumin has increase, comparison with control group.

Low serum urea concentrations have been recognized previously in association with liver failure and have been suggested to indicate reduced hepatic synthesis of urea from ammonia. The decreased serum urea is associated with more severe hepatopathies and has prognostic relevance. on the other hand, it may be absorbed by the hydatid cyst since (65) detected a large number of carbohydrate, Aspartate aminotransferase (AST) was also measured showed an increase among the positive group compared to the negative control group; a result agreed with (64) who found that the liver infection with cestoda tapeworms led to hepatocyte destruction and enzyme release, therefore, the concentration of AST increased.

The values of these indicators vary; several indicators are in normal limits and some change. Determination of the activity of some enzymes in serum gives us a more complete picture of activity and functional status of the liver. We have selected some enzymes, but also can serve as biological markers with diagnostic value. It has been reported that host biochemistry shows some alterations from normal physiology in animals with



parasitic infections (66,67,62). In studies conducted on parasite infested animals (68,69), changes in total serum protein levels, and the levels of certain vitamins, enzymes, and minerals were reported, while the parasitic diseases were reported to make infested animals susceptible to vitamin and study conducted by (66) found that there were some increases in some biochemical parameters including total protein, globulin, amylase, chlor, and vitamin B12 levels, and some decreases in albumin, magnesium, and phosphorus levels of sheep diagnosed with cyst hydatid (66). In the physiopathology of the liver, AST, ALT, ALP, GGT, CPK, bilirubin and albumin, globulin protein tests are important parameters (70). Increased AST activity in the serum is a sensitive marker of liver damage (71,72) report that the ALP and AST activities of sheep infested with endoparasites are higher compared to healthy animals and that the increase is caused by the necrosis or cholestasis of the hepatic cells. It is believed that the increase in AST activity and bilirubin levels might be due to hepatocytes, while the increase in total protein and necrosis and/or cholestasis of the globulin levels might be attributed to the deterioration of the protein metabolism.

TREATMENT

Medication

Small, superficial cysts in one location may respond to antiparasitic drugs called benzimidazoles. Benzimidazoles destroy parasites and shrink cysts. Your healthcare provider may combine medication with aspiration or surgery.

PAIR

(puncture, aspiration, injection, re-aspiration): The PAIR technique uses a needle or catheter to drain the cyst. A parasite-fighting chemical is injected into the cyst before it's drained again. Your healthcare provider repeats the process until the cyst is completely emptied.

SURGERY

Furthermore, despite the availability of novel therapeutic options, surgery remains the most prevalent treatment option for this condition, resulting in significant economic and health consequences for nations (41), hydatid cysts, according to certain research, have distinct clinical-radiological features and treatment approaches, clinicians frequently struggle to make a solid diagnosis for this condition (42) Parasitic infection seem to cause increased hepatic enzyme activity and decreased trace element levels by causing liver damage (43,44)

PREVENTION

Control measures discussed above are directed toward prevention of infection. In relation to such measures, the prevention of human infection by *E. granulosus* eggs is based primarily on health education. It is important to determine the personal, socioeconomic, and environmental risk factors involved in the transmission of human hydatidosis in each country or region. One risk is the lack of potable water and the use of rural water tanks that collect rainwater from the ground surface and roofs. All personnel handling dogs in endemic areas should be aware of the health risk of acquiring hydatidosis and must take safety precautions. At a personal level, the prevention is mainly based on simple rules of personal hygiene, such as avoiding contact with unknown dogs that may be

infected, washing hands before eating, and washing vegetables before eating them raw. Dog owners should treat the dogs regularly with praziquantel and avoid feeding them with raw offal potentially infected with cysts.

CONCLUSION

The number of eosinophils shows interest because it can be used routinely as a biological indicator for establishing a preliminary diagnosis which is likely accurate for parasitic. Hematological indicators with diagnostic value, resulted HB, PCV, eosinophils. Their evaluation, can be used as a routine screening test from districts laboratories determining preliminary diagnosis in sheep and mice infested of liver parasites. The change of hepatic enzymes level serves to monitor the progress of parasitic infection in animals and as a sensitive diagnostic aid in field infections. Albumin, BUN, creatinine and total bilirubin are reliable indicators of the stage and severity of parasitic liver naturally infested sheep and in doing so, and constitute an important diagnostic tool in determining of the official diagnosis and an efficient treatment process. The results of the study indicate that the determination of the changes in AST, total protein, total bilirubin levels of the animals infected with cystic echinococcosis could be used as assistive laboratory analyzes to determine the physio pathological changes in their livers and the prognosis of the diseased animals.

REFERENCES

1. Lötsch, F; Budke, C. M; Auer, H., Kaczierek, K; Waneck, F Lagler, H. and Ramharter, M. (2019). Evaluation of direct costs associated with alveolar and cystic Echinococcosis in Austria. *PLoS Neglected Tropical Diseases.*, 13(1): 7110p
2. Higueta, N. I. A; Brunetti, E., and McCloskey, C. (2016). Cystic echinococcosis. *Journal of clinical microbiology*, 54(3): 518-523
3. Yildiz K, Gurcan S 2003. Prevalence of hydatidosis and fertility of hydatid cysts in sheep in Kirikkale, Turkey. *Acta Veterinaria Hungarica*, 51 (2): 181- 187
4. Hakverdi S, Culha G, Canda MS, Yaldiz M, Altintas S 2008. Problem of Cystic Echinococcosis in Hatay. *Turkiye Parazit Derg*, 32: 340-342.
5. Demir P, Mor N 2011. Seasonal Distribution and Economic Importance of Cystic Echinococcosis in Cattle Slaughtered at Kars Municipal Abattoir, Turkey. *Turkiye Parazit Derg*, 35: 185-188.
6. Nunnari G, Pinzone MR, Gruttadauria S, et al. Hepatic echinococcosis: clinical and therapeutic aspects. *World J Gastroenterol*. 2012; 18:1448-1458.
7. Tiaoying, L.; Jiamin, Q.; Wen, Y.; Craig, P.S.; Xingwang, C. and Schantz, O.M. (2005). Echinococcosis in Tibetan populations, western Sichuan province, china. *Emerg. Infect. Dis.*, 11: 1866 – 1873.
8. Romig, T. (2003). *Epidemiology of echinococcosis, Langenbecks Arch. Surg.* 388:209-17.
9. Parija, S.C. and Sheela, D. C. (1999). Current concepts in the diagnosis of cystic echinococcosis in humans and Livestocks and intestinal echinococcosis in canine hosts. *J. vet. Parasit.*, 13: 93-102.
10. Radostits O. M., Gay C.C., Hinchcliff K. W., Constable P. D. (2009). *Veterinary Medicine A textbook of the diseases of cattle, horses, sheep, pigs and goats Tenth Edition*, 2009.



11. Pedrosa I, Saiz A, Arrazola J, Ferreiros J, Pedrosa CS. Hydatid disease: radiologic and pathologic features and complications. *Radiographics*. 2000; 20:795-817.
12. Eckert J, Deplazes P. Biological, epidemiological, and clinical aspects of echinococcosis, a zoonosis of increasing concern. *Clin Microbiol Rev*. 2004; 17:107-135.
13. Yildiz K, Gurcan S 2003. Prevalence of hydatidosis and fertility of hydatid cysts in sheep in Kirikkale, Turkey. *Acta Veterinaria Hungarica*, 51 (2): 181- 187.
14. Hakoverdi S, Culha G, Canda MS, Yaldiz M, Altintas S 2008. Problem of Cystic Echinococcosis in Hatay. *Turkiye Parazit Derg*, 32: 340-342.
15. Demir P, Mor N 2011. Seasonal Distribution and Economic Importance of Cystic Echinococcosis in Cattle Slaughtered at Kars Municipal Abattoir, Turkey. *Turkiye Parazit Derg*, 35: 185-188.
16. Hayajneh, F. M. F; Althomali, A. M. H; and Nasr, A. T. (2014) Prevalence and characterization of hydatidosis in animals slaughtered at Al-Taif abattoir, Kingdom of Saudi Arabia. *Open Journal of Animal Sciences*, 4(01), 38- 41
17. Ruptured hydatid cyst of lung. *Current Pediatric Research. Scientific Publishers of India.*, 16 (2): 156-158
18. Khalifa, R; Nasser, F; Elsetouhy, A; and Farag, I. (2016). Hydatid cyst of the neck. A case report and literature review. *Egyptian Journal of Ear, Nose, Throat and Allied Sciences*, 17(2): 103-105
19. Abdulhameed, M. F; Robertson, L. D., Al-Azizz, S. A., and Habib I (2019). Neglected zoonoses and the missing opportunities for one health education: The case of cystic echinococcosis among operated patients in Basrah, Southern Iraq *Diseases Surgically Journal*, 7(4): 1- 10
20. Thompson, R. C. A. (2017). Biology and systematics of *Echinococcus* in *Advances in parasitology*. Academic Press 95: 65-109
21. Barnes, T.; Deplazes, P.; Gottstein, B. (2012). Challenges for diagnosis and control of cystic hydatid disease. *Acta Trop.*, 123:1-7
22. Dakkak, A (2010). Echinococcosis / hydatidosis: a severe threat in Mediterranean countries. *Veterinary Parasitology.*, 174(1-2): 2-11
23. Luma Abd Almunim Baker, Shaymaa Zuhir Jalal Aldin. (2022). Association of some biochemical parameters and blood pressure among males with hypertension in the camps of Nineveh province Iraq. *Journal of Population Therapeutics and Clinical Pharmacology*, 29(04), 167-176.
24. Hamdani A, Khawar R, Rubab H. Aggressive Behavior and Coping Strategies among Cardiac Patients with Type A and Type B Behavior Pattern. *Pak Heart J*. 2022;55(03):242-246.
25. Ashraf T, Ishaq M. Coronary Artery Disease and Depression: A Missing Pillar in Management. *Pak Heart J*. 2022;55(03):205- 206.
26. Ashiq S, Ashiq K, Sabar MF. The Complex Relationship of Paraoxonase Gene Polymorphisms with Coronary Artery Disease and Lipid Metabolism. *Pak Heart J*. 2022;55(03):307-
27. Gokcen A, Sahin T, Camkerten I 2006. A primary case of Cardiac cyst hydatid in a calf, in Turkey. *Yüzüncü Yıl Üniversitesi Veteriner Fakültesi Dergisi*, 17 (1): 33-35.
28. Karaman U, Miman O, Kara M, Gicik Y, Aycan OM, Atambay M 2005. Hydatid Cyst Prevalence in the Region of Kars. *Türkiye Parazit Derg*, 29 (4): 238- 240.
29. Ayaz E, Ertekin A, Ozdal N, Tas Z 2006. Some Biochemical Parameters in Sheep Infected with Endoparasites (*Fasciola* spp., *Dicrocoelium dendriticum*, *Hydatid* cysts, *Trichostrongylidae* and *Protostrongylidae*). *Turkiye Parazit Derg*, 30 (4): 57-61
30. P. Moro and P. M. Schantz, "Echinococcosis: a review," *International journal of Infectious diseases*, vol. 13, no. 2, pp. 125-133, 2009.
31. D. Ghartimagar, A. Ghosh, M. K. Shrestha, O. P. Talwar, and B. Sathian, "14 years' hospital based study on clinical and morphological spectrum of hydatid disease," *Journal of Nepal Medical Association*, vol. 52, no. 190, pp. 349-353, 2013.
32. R. Morar and C. Feldman, "Pulmonary echinococcosis," *European Respiratory Journal*, vol. 21, no. 6, pp. 1069-1077, 2003.
33. H. Salama, M. F. Abdel-Wahab, and G. T. Strickland, "Diagnosis and treatment of hepatic hydatid cysts with the aid of echo-guided percutaneous cyst puncture," *Clinical infectious diseases*, vol. 21, no. 6, pp. 1372-1376, 1995.
34. P. L. Moro, *Clinical manifestations and diagnosis of echinococcosis*, vol. 1, 2016.
35. Anand S., Col L., Rajagopalan S., Brig B. and Mohan R. Management of liver hydatid cysts - Current perspectives, *Med J Armed Forces India*, 2017;73(1):104.
36. Heymann D. (2015). *Control of Communicable Diseases Manual*. (20th edition). Washington, DC: American Public Health Association, pp178 - 183.
37. Ibrahim Z. A. Attenuation of human hydatid cyst protoscolices viability by 1-Hydroxyphenazin (1-HP) pigment: In vitro and in vivo study, *Journal of the Faculty of Medicine Baghdad*, 2010;52(3): 338-41.
38. W. S. Kammerer and P. M. Schantz, "Echinococcal disease," *Infectious disease clinics of North America*, vol. 7, no. 3, pp. 605-618, 1993.
39. P. Moro and P. M. Schantz, "Echinococcosis: a review," *International journal of Infectious diseases*, vol. 13, no. 2, pp. 125-133, 2009.
40. Akkapulu N, Aytac HO, Arer IM, Kus M, Yabanoglu H. Incidence and risk factors of biliary fistulation from a hepatic hydatid cyst in clinically asymptomatic patients. *Tropical doctor*. 2018;48(1):20-4.
41. Ahmed J, Arif TB, Tahir F, Malik F, Parkash O. Hydatid cyst of spleen presenting with vague symptoms: a diagnostic conundrum. *Cureus*.
42. Khamees Abed S, Najim TM, Shahooth MA. Isolation of Bacteria Associated with Hepatic Hydatid Cyst of Iraqi Sheep. *European Journal of Molecular & Clinical Medicine*. 2020;7(11):972-6.
43. Vengust G, Klinkon M, Bidovec A, Vengust A 2003. *Fasciola hepatica*: effect on blood constituents and liver minerals in fallow deer (*Dama dama*). *Vet Parasitol*, 112: 51-61.
44. Sahin T, Akgul Y 2006. Investigation of Some Trace Element Levels and Biochemical Parameters in Sheep with Endoparasite. *YYÜ Sağlık Bilimleri Dergisi*, 9 (1): 100-106.
45. Humairy, A. KH. (2010). Evaluation of the Activity of *Datura stramonium* Seeds Extracts on Growth and Development of *Hydatid* Cysts for *Echinococcus granulosus* in White Mice Balb/c (Therapeutic, Histologic and Immunologic Study). Ph. D. Thesis. College of Science, Kufa University
46. Al - Mobarak, Z.A. (2006). Effect of peel extracts of *Punica* fruits on echinococcosis of *Echinococcus granulosus*. M. Sc. Thesis, College of Education., Kufa University, (In Arabic).



47. Al-Nasiri, F.S. (2006). *Biological and immunological study of hydatid cyst formation in albino mice*. Ph.D. Thesis, College of Education, University of Baghdad, (In Arabic).
48. Moraitaki, P.; Roussi, P.; Emmanouil, Y. Pechlivanidou, R.; Bitsakou, C.; Konstantinou, M. and Marosis, K. (2010). Pulmonary echinococcosis presenting as a pulmonary mass with fever and haemoptysis; a case report. *Pneumon. J.* 23, (2): 180-183
49. Nguyen, D.T. and Diamond, L.W. (2000). *Diagnostic hematology*. 1st ed. Butter Worth Heinemann, Oxford. P.96 - 98.
50. Hindson J.C., Winter A. C., *Manual of Sheep Diseases*, Second Edition, 2002.
51. Kramer, J.W. *Normal haematology of cattle, sheep, and goat*. In: FELDMAN, B.F.; ZINKL, J.G.; JAIN, N.C. (Eds). *Schalm's veterinary haematology*. 5.ed. Philadelphia: Lippincott Williams & Wilkins, 2000. p.1075-1084.
52. Radostits O. M., Gay C.C., Hinchcliff K. W., Constable P. D. (2009). *Veterinary Medicine A textbook of the diseases cattle, horses, sheep, pigs and goats Tenth Edition*, 2009.
53. Mekroud1 A., Chauvin A. 2. and Rondelaud D. 3 *. *Variations of biological indicators as highly presumptive markers fasciolosis in experimentally-infected sheep*. *Revue Méd. Vét., for 2007*, 158, 8-9, 437-441.
54. Sahin T, Akgul Y 2006. *Investigation of Some Trace Element Levels and Biochemical Parameters in Sheep with Endoparasite*. *YYÜ Sağlık Bilimleri Dergisi*, 9 (1): 100-106.
55. Mert N, Kozat S, Ekin S, Gunduz H, Denizhan V 2003. *Doğal kronik Fasciolasisli koyunlarda serum sialik asit ve serum lipid-bağlı sialik asit düzeyleri*. 13. Ulusal Parazitoloji Kongresi, 8-12 Eylül 2003, Konya, Turkey.
56. Tanritanir P, Özdal N, Ragbetli C, Yörük I, Ceylan E, Deger S 2009. *Some biochemical parameters and vitamins levels in the hair goats naturally mix infested with endo-ectoparasites (lice (linognathus africanus) and Trichostrongylidae sp.)*. *J Anim Vet Adv*, 9: 1707-1711.
57. Uslu U, Akalin PP, Şahin EY, Altunok V 2011. *Some biochemical parameters in yearling sheep naturally infected with Coenurus Cerebralis*. *Fresenius Environmental Bulletin*, 20(2): 411-414.
58. Kam K 2016. *Investigation of mineral content of the serum and cyst fluid in sheep with hydatid cysts*. (Master thesis), Yuzuncu Yil University, Institute of Health Sciences Department of Biochemistry, Van, Turkey.
59. Vengust G, Klinkon M, Bidovec A, Vengust A 2003. *Fasciola hepatica: effect on blood constituents and liver minerals in fallow deer (Dama dama)*. *Vet Parasitol*, 112: 51-61
60. Karagul H, Altıntaş A, Fidancı UR, Sel T 2000. *Klinik Biyokimya*. 1st ed. Ankara, Turkey: Medisan Yayınevi.
61. Tennant BC, Center SA 2008. *Chapter 13: hepatic function*. In *Clinical Biochemistry of Domestic Animals*. sixth ed. In: Kaneko, J.J., Harvey, J.W., Bruss, M.L. (Eds.), Academic Press, New York, USA, pp. 379-412.
62. Shaafie IA, Khan AH, Rambabu K. *Biochemical profiles of hydatid cyst fluids of Echinococcus granulosus of human and animal origin in Libya*. *J Helminthol*. 1999; 73:255-258.
63. Caremani M, Lapini L, Caremani D, Occhini U. *Sonographic diagnosis of hydatidosis: the sign of the cyst walls*. *Eur J Ultrasound*. 2003; 16:217-223.
64. Abdullah, B.H. (2007). *Some Helminths Parasitized on Stray Cats Felis catus L. in Basrah and Epizootiological and Biological Studies on Taenia taeniaeformis Batsch, 1786*. Ph.D. Thesis, College of Education, University of Basrah, (In Arabic).
65. Ozen, N.; Celik, C.; Ozkan, K.; Malazgirt, Z.; Isimer, A. and Sayal A. (1992). *Trace elements in hydatid disease*. *Journal of Trace Elements and Electrolytes in Health and Disease*. 6: 67-70.
66. Ayaz E, Ertekin A, Ozdal N, Tas Z 2006. *Some Biochemical Parameters in Sheep Infected with Endoparasites (Fasciola spp., Dicrocoelium dendriticum, Hydatid cysts, Trichostrongylidae and Protostrongylidae)*. *Türkiye Parazit Derg*, 30 (4): 57-61.
67. Deger Y, Gul A, Bildik A, Dede S, Yur F, Deger S 1997. *Parazitli köpeklerin bazı kan parametreleri ile plazma vitamin C düzeylerinde görülen değişiklikler*. *Türkiye Parazit Derg*, 21: 195-198.
68. Tanyuksel M, Sayal A, Aydın A 1995. *Paraziter hastalıklarda eser element düzeyleri*. *Türkiye Parazit Derg*, 19: 315-321.
69. Mert N, Kozat S, Ekin S, Gunduz H, Denizhan V 2003. *Doğal kronik Fasciolasisli koyunlarda serum sialik asit ve serum lipid-bağlı sialik asit düzeyleri*. 13. Ulusal Parazitoloji Kongresi, 8-12 Eylül 2003, Konya, Turkey.
70. Comba B, Mert H, Comba A, Mis L, Mert N 2017. *The Some Hematological and Biochemical Parameters in Karakul and Norduz Sheep*. *Van Vet J*, 28 (3): 137-140.
71. Meyer DJ, Harvey JW 1998. *Veterinary Laboratory Medicine. Interpretation and Diagnosis*. 2nd ed., W. B. Saunders Company. Philadelphia, London, Toronto, Montreal, Sydney, Tokyo. pp. 157-187.
72. Yuksek N, Altug N, Gul A 2007. *Therapeutic effect of the combination of trichlobendazole and leвамisol in sheep with endoparasite infection*. *YYU Vet FakDerg*, 18: 19-24.