ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188

NEEDLE LITTER DECOMPOSITION AND NUTRIENT RELEASE IN SUB-TROPICAL PINE FOREST OF MANIPUR, NORTH-EAST, INDIA

Dr. Khulem Ujala Devi
Assistant Professor, Department of Botany, Liberal College, Imphal, Manipur

Article DOI: https://doi.org/10.36713/epra25739
DOI No: 10.36713/epra25739

ABSTRACT

*A faster rate of needle litter decomposition of Pinus kesiya occurred in forest site I (undisturbed) with losing 98% of the initial mass in comparison with 96% of mass loss for forest site II (disturbed) over 24 months. Decomposition rate shows seasonal changes with the maximum rate in rainy season and minimum in dry winter season. Needle litter of forest site I exhibited higher initial concentration of nitrogen, cellulose and lower lignin and lignin to nitrogen ratio than forest site II. The rate of decomposition was correlated with soil moisture (r = 0.51**, 0.51**); soil temperature (r = 0.55**, 0.52**) than rainfall and mean air temperature in both forest sites. Chemical composition like initial nitrogen, lignin, lignin to nitrogen ratio and cellulose were significantly correlated with remaining biomass. (r = 0.78, 0.95, 0.95 and 0.81) in forest site I and (r = 0.62, 0.90, 0.91 and 0.75) in forest site II at the level of P < 0.01 in both forest sites. The decay constant were 0.594 and 0.582 in forest site I and site II respectively. During decomposition, concentrations of nitrogen and phosphorous were found to increase whereas potassium and sodium exhibited decreasing pattern from the original concentration in both the forest sites. The average concentration of nutrients in both forest sites was in the order N > K > P > Na. The release of nutrients (N P K and Na) based on the amount of element remaining over two years in forest site I and forest site II was in the order K>Na > P > N.*

KEY WORDS : *Pinus Kesiya, Needle Litter, Decomposition, Nutrient Concentration, Nutrient Release, Lignin, Cellulose And North-East India (Manipur).*

1. INTRODUCTION

Decomposition of litter is an important process in most terrestrial ecosystems because of its role in regulating building up of soil organic matter, releasing of nutrients for plant growth and influencing carbon flux from soil (Wardle *et al.,* 2003; Ganjeguente *et al.,* 2004). The decomposition of needle litter can be divided into two phases. In the first phase, decomposition rate is regulated by the concentration of nutrients in the litter and lignin decomposition rate in the second phase (Berg and Staaf, 1980). Leaf litter, which accounts for a large proportion of dead organic matter (~70%) (Costa et al. 2020), is the major source of nutrients and energy for heterotrophic communities in forest soil (García-Palacios et al. 2016a; Jia et al. 2018)

Decomposition and nutrient release rates are influenced by its resource quality, the activity and composition of soil organism and the physical micro-environment (Meentemeyer, 1978; Seastedt *et al.,* 1983; Finzi *et al.,* 2001; Gonzalez and Sealedt, 2001; Bradford *et al.,* 2002; Arunachalam & Singh, 2004; Goma & Bernhard, 2006). Resource quality includes N concentration, C - N ratio, lignin concentration, lignin to nitrogen ratio, soluble polyphenols and polyphenol - nitrogen ratios (Berg and Staaf, 1991; Mellilo *et al.,* 1982; Palm and Sanchez, 1991; Constantinides and Fownes, 1994; Mtambanengwe and Kirchmann, 1995). Under the influence of the prevalling elimatic environment, different litter have their own specific rate of decomposition which govern the rate at which nutrients are released from litter. The dynamics of the nutrients during decomposition of litter is complex because the nutrients often occur in different forms and are subject to various transformation.

High contents of lignin have a rate reducing influence on litter decomposition both in the early and later stages particularly when associated with high N content (Berg and Ekbohm, 1991; Couteaux *et al.,* 1995). Never the less a high nitrogen content generally regulates the early stages by enhaching the growth of microorganisms that degrade labile compounds and repress the form action of lignolytic enzymes (Keyser *et al.,* 1978). The role in cellulose and lignin is regulating factors of litter decomposition rate in temperate, boreal coniferous and deciduous forest (Mc Claugherty and Berg, 1987; Berg *et al.,* 1996; Couteaux *et. al.,* 1998).

Objective : To Evaluate

1. the rate of needle litter decomposition,
2. the release of nutrients (N, P, K & Na) and
3. the release rate of lignin and cellulose in bimonthly intervals of pine forest site I and forest site II.

2.MATERIALS AND METHODS

2.1. Site Description

The present investigation was conducted in two sub-tropical pine forest ecosystems of Imphal west district, Manipur (India) which falls under Eastern Himalayan region. The pine forest of Engel Ching (Engel Hill) is an undisturbed site (Forest site I) located near Tendongyan village and it falls at $24^0 55'$N latitude

ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188

and 93^{0}55^{l}E longitude at an altitude of 895m from the mean sea level. The another pine forest is disturbed site (forest site II) extending on the Langol hill ranges near the Khamaran village which lies at 24^{0}52^{l}N latitude and 93^{0}54^{l}E longitude at an altitude of 960m from the mean sea level. The climate is monsoonic and the average annual rainfall is 1429.79mm. Soil type is red sand stone mixed with gravel. Soil P^H ranged from 5.11 to 5.47. The mean maximum temperature ranged between 26.0 to 26.20^{0}C and mean minimum temperature ranged from 12.60^{0}C to 14.90^{0}C.

2.2. Field Experiment

Litterbag method was followed to estimated the rate of needle litter decomposition of *Pinus kesiya* in the forest floor (Gilbert and Bocock, 1962). Litter bags of nylon net with a mesh size of 1mm and an over all size of 10cm × 10cm were used. A mesh size of 1mm was sufficient for movement of micro arthropods which are the predominant litter feeders in the forests (Sharma *et al.,* 1984). Five grams of needle leaves were kept in nylon litterbags. The litter bags were placed on the forest floor in three different zones (base, middle, top) of hill and 36 litterbags were used for each zones and a total of 108 (12 × 3 × 3) literbags were placed in each forest site and clip with nail. However care being taken so that the samples were not disturbed on the forest floor. Nine litter bags were collected from each forest sites at bimonthly intervals from the month of October 2004 to August 2006. The recovered litter bags were brought to the laboratory. The residual materials from the litter bags were washed under a fine jet of water using a 1000 *m* mesh screen to remove all the adhering soil particles and dried in hot air oven at 80^{0}C to constant weight and then ground for chemical analysis. Soil temperature and soil moisture were measured using thermometer and moisture meter. Mean air temperature and rainfall was recorded from the Meteorological Department, Tulihal Airport, Imphal, Manipur (India).

2.3. Data Analysis

The decomposition constant (K) of the decomposing needle litter for the entire investigation period was calculated following exponential decay model (Olson, 1963). e^{-kt} = DM / DMo. Further, the time required for 50% (t_{50}) and 95% (t_{95}) decay was calculated as t_{50} = 0.693 / K and t_{95} = 3/K respectively. Absolute amount remaining in the needle liter after a given period of time was determined by using the following equation. Percent absolute nutrient remaining = C/Co × DM/DMo × 100 where, DMo is the original mass of dry matter, DM is the mass of dry matter after a given period, K is annual decomposition (yr^{-1}), t is the time (yr), Co is the original constant of element in needle litter and C is the concentration after a given period.

2.4. Chemical Analysis

The estimation of nitrogen and phosphorus were determined following the method given by Anderson and Ingram (1993) using Calorimeter and Potassium and sodium by Flame photometer. Lignin content of needle leaves were estimated according to Peach and Tracy (1955). Cellulose of the needle were estimated after Up-degraffs D.M. Method 1969.

3.RESULT

3.1. Litter Decomposition

The percentage of original biomass remaining of needle litter decomposition in different months in pine forest site I and site II are given in fig. 1. Higher rate of litter decomposition was recorded during rainy months from August '04 through October '04 and from June '05 through October '05 and lower rate during dry winter months. The rate of needle litter decomposition was recorded to be 53% and 49% after the completion of one year (October 2004 to August 2005) of decomposition study in forest site I and forest site II. However, at the termination of the experiment 2% and 4% of original biomass was remained in forest site I and forest site II respectively.



Fig. 1. Bimonthly variation of remaining biomass in the needle litter decomposition in pine forest site I and forest site II.

ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188

3.2. Initial Litter Chemistry

The initial nitrogen, lignin, lignin to nitrogen ratio, cellulose of the initial litter material are presented in table 1.

Table 1 : Initial N, L. L to N ratio and cellulose in the pine forest site I and site II.

Forest	IN	IL	IL/IN	IC
Site I	0.60	27	45	20
Site II	0.59	32	54	17

IN - Initial nitrogen concentration (%), IL - Initial lignin ,
IL/IN - Initial lignin to nitrogen ratio, IC - Initial cellulose (%)

3.3. Correlations between variables

The rate of decomposition in different months was significantly correlated with the soil moisture (r = 0.51**, 0.51**); soil temperature (r = 0.55**, 0.52**) where as low correlation and significant was evident with mean air temperature (r = 0.10*, 0.16*) in forest site I and forest site II (table 2). However, rainfall showed low correlation and significant (r = 0.42*) and significant correlation (r = 0.51*) in forest site I and forest site II respectively. The variability of SM, ST, RF and MAT in forest site I and forest site II are showed in table 2.

The remaining biomass (%) showed positive correlation and significant with initial nitrogen, initial lignin, initial lignin to nitrogen ratio and initial cellulose (r = 0.78, 0.95, 0.95 & 0.87) and (r = 0.62, 0.90, 0.91 & 0.75) at the level of P < 0.01 in forest site I and site II respectively. The variability observed by initial nitrogen, initial lignin, initial lignin to nitrogen and initial cellulose are 61%, 90%, 90% and 76% in forest site I and 38%, 81%, 83% and 56% in forest site II.

Table 2 : Correlation between Rate of weight loss and abiotic variables in forest site I and site II (n = 12)

Parameter	Forest Site I		Forest Site II	
	R	Variability(%)	R	Variability(%)
Siol Moisture (SM)	0.51**	26	0.51**	26
Soil Temp.(ST)	0.55**	30	0.52**	27
Rainfall(RF)	0.42*	18	0.51*	26
Mean air Temp.(MAT)	0.10*	1	0.16*	3

* shows significant at P<0.05 level
** shows significant of P<0.01 level.

3.4. Decomposition Constant

Decomposition constant (K), as well as half time (t_{50}) and time to 95% loss of dry weight (t_{95}) are presented in table 3. The annual decomposition constant (K) was recorded to be 0.591 and 0.582 in forest site I and forest site II respectively.

Tables 3 : Rate of needle litter decomposition, turnover rate ,decomposition rate and constant (K)during litter decomposition in pine forest site I and forest site II.

Forest	Rate of decomposition	Turnover rate	Decomposition (% per day)	constant (K)
Site I	0.27	1.173	5.069	0.591
Site II	0.26	1.190	5.147	0.582

3.5. Nutrient dynamics in the decomposing needle litter

Changes in relative concentration and absolute amount or percent remaining of nitrogen, phosphorus, potassium and sodium in the residual needle litter of forest site I and site II are presented in Fig. 2. Nitrogen concentration in decomposing litter increased significantly from the original concentration (0.60%) and (0.59%) in forest site I and forest site II respectively (Fig. 2.A). However it exhibited some fluctuation followed by a rapid increased to 1.50% in forest site I and 1.17% in forest site II after 24 months elapsed.

A net immobilization of nitrogen above 100% of the initial amount was exhibited from the beginning upto December 2004 in forest site I and June 2005 in forest site II (Fig. 2.B). However nitrogen mineralization continued till the end of study and remained only 5% and 8% of original N mass in forest site I and site II respectively.

Phosphorous concentration slightly reduced then the original concentration (0.12% and 0.13%) in early stages then increased by 0.16% and 0.15% in both forest site I and site II respectively after 24 months elapsed (Fig. 2.C). There was a great released of phosphorous in the first stage of month (October2004) 26% in forest site I and 23% in forest site II (Fig. 2.D.). The pattern of P flux in both the forests sites was irregular and remained 3% and 5% after 24 months.

The concentration of potassium declined rapidly to 0.08% and 0.09% in the first month (October 2004) in both the forest sites then slowly increased by 0.20% (August, 2005) and thereafter fluctuated (Fig. 2E). Potassium was released rapidly in the initial month of October 2004 (78% and 72%) followed by slowly increased upto June 2005 but not exceed from the initial amount then released later on in both forest sites (2.F). After 24 months only 1% and 2% of K was remained in forest site I and site II respectively.

ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188

The initial concentration of sodium in both the forest sites was 0.04% and decreasing concentration started from February 2005 (0.03%) and at the end of the assay recorded 0.02% in both forest sites (2.G). In forest site I, sodium exhibited gradual net release from the beginning till the end of assay whereas in forest site II observed an initial net release phase upto December 2004 in which 35% of Na was lost and followed by 84% of initial Na content (February'05) then release continously. Only 1% and 4% of Na was remained in forest site I and site II respectively (2.H).

4. Lignin and Cellulose Concentration During Decomposition

The lignin concentration during decomposition of needle litter increases from the original concentration (Table 1. Fig. 3 A). There is increasing trend from the initial month till the end of the study was 45% and 50% in forest site I and site II respectively. However, it was recorded some fluctuation from the month of February to June 2005 in forest sites II. Cellulose concentration in contrast, exhibited lower then the original concentration (Table 1. Fig. 3 B) in both forest sites. After 2 years of decomposition cellulose concentration was 5% and 6% in forest site I and site II respectively.




Fig. 2. Changes of N (A), P (C), K(E) and Na (G) concentration and % remaining N (B), P (D), K (F) and Na (H) in decomposing needle litter of forest site I and site II.

ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188




Fig. 3 Variation of lignin (3.A), celloluse (3.B) concentration in decomposing needle litter of forest site I and site II.

4.DISCUSSION

The rate of needle litter decomposition in bimonthly intervals shows different variations. The higher rate of decomposition during rainy season then in winter season in both forest sites may be due to high percentage of soil moisture and favourable soil temperature which enhances the activity of microorganisms and water soluble fractions in litter leached out more in wet summer season. Marchante et.al,2019 and Akoto et.al,2022 also reported higher rate of decomposition in rainy season and lower in winter season.

In both forest site I and site II the rate of weight loss was more during the first year (53% and 49%), then the second year (45% & 47%) and the value was comparable with the pine forest of Central Himalaya (Upadhyay and Singh, 1989), in *Casuarina equisetifolia* plantation in Orissa, India. (Misra and Nisanka, 1997) and *Populus deltoides* in Himachal Pradesh, India (Kaushal *et al.,* 2006). The total rate of needle litter decomposition of forest site I (98%) is higher than forest site II (96%) after 24 months. It may be due to the higher initial level of N, lower initial lignin, higher initial cellulose and lower initial lignin to nitrogen ratio of forest site I than forest site II (Table 1). Several workers reported that high initial level of N, low initial level of lignin, higher rate of initial cellulose and low initial lignin to nitrogen ratio resulted in the high rate of decomposition (Melillo *et al.,* 1982; Webderburn and Carter, 1999; Sarjubala, 2002; Bijayalaxmi, 2005).

In the needle litter decomposition significant positive correlation was observed with soil moisture (r=0.51**, 0.51**) and soil temperature (r=0.55**, 0.52**) than rainfall and mean air temperature in both forest site I and site II (Table 2). The maximum variability was recorded in soil temperature (30%, 27%) and soil moisture (26%). It is obvious that soil temperature and soil moisture are highly correlated with litter decomposition.

Positive and significant of remaining biomass (%) was observed with initial nitrogen, initial lignin, initial lignin to nitrogen ratio and initial cellulose (r=0.78, 0.95, 0.95 & 0.87) and (r = 0.62, 0.90; 0.91 & 0.75) at P<0.01 in forest site I and

ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188

site II respectively. The variability values by initial lignin (90% and 81%), initial lignin to nitrogen (90% and 83%) shows maximum percentage than nitrogen and cellulose. It has been found that IN and IL / IN was more correlated with remaining biomass in both forest sites. David *et al.,* (1998) showed correlation of remaining mass with initial lignin and nitrogen ratio ($r = 0.74$, $P < 0.01$) in *Pinus sylvestris* in Grizedala forest, Cumbria, U.K. Upadhyay *et al.,* 1989 also reported a significant correlation of remaining biomass with lignin ($r=0.80$, $P<0.01$), lignin to nitrogen ratio ($r=0.58$, $P<0.05$) in Central Himalayan forest of India.

The rate of decomposition constant 0.59 of forest site I is little higher than forest site II (0.58). Higher rate of decomposition constant indicates the higher rate of decomposition. The present value is comparable with pine forest at Meghalaya, Shillong (0.74) by Das and Ramakrishnan, 1985; Pine forest of Europe (0.32-0.56) by Kurz *et.al.,*2005 and pine forest in U.S.A. (0.27 _ 0.54) by Cusack *et al.,* 2009 but higher than that of *Pinus densiflora* (0.21) by Namgung *et al.,* 2008 at Korea.

Nitrogen concentration in needle litter decomposition increased from the original concentration in both forest sites. Such an increased could be due to nitrogen mineralization both microbial and non microbial and deposition of nitrogen from atmosphere (Jorgenson and Myer, 1990; Kocky & Wilson, 1997). Change in the absolute amount of an element during decomposition (net immobilization or net release) is a function of both mass loss and change in residual litter. A value greater than 100% indicated immobilization of the element whereas value less than 100% indicated the release of the element from the litter.

N immobilization occurred from the beginning through December 2004 in forest site I and June 2005 in forest site II, thereafter occurred N mineralization. N immobilization in the early stages is related with increased concentration of nitrogen during decomposition in both forest sites (Fig. 2 A, B). Mun (2009) reported increased N concentration during decomposition resulted N immobilization in the early period in Oak forest, at Korea. Pankaj et.al, 2022 also reported increased N and P concentration in decomposing litter. The release of N (95%) in forest site I was found higher than forest site II (92%). It may be due to the increased of initial nitrogen concentration in forest site I (table 1). Cusack (2009) reported N release was higher in species with a high initial N concentration in pine forest at U.S.A.

Phosphorus concentration slightly reduced then the original concentration (0.12 & 0.13%) in early stages and then increased by 0.16% and 0.15% in forest site I and site II respectively after 24 months elapsed. Increased P concentration may be attributed to input throughfall and biological translocation (by fungi) from deeper soil layers (Mc Brayer and Cromark, 1980; Xu *et al.,* 2004). The rapid phosphorus loss observed in the first stage (October 2004) (Fig. 2.D) was probably a consequence of leaching of soluble P containing compounds. Many workers reported the leaching of P in the early stages of decomposition (Blair, 1988; Tripathi and Singh, 1992; Mun, 2009). The initial leaching phase was followed by some degree of P increased but did not exceed the initial amount in both the forest sites. It may be due to microbial immobilization (Bockheim *et al.,*1991).

The concentration of potassium declined rapidly from the original concentration in both forest sites. Absolute amount of K was also released rapidly in the initial months (October 2004) from the original K mass in which 78% and 72% of K was lost and then slowly releases. It can be observed that decreased rate of K concentration related with rapid release of K mass. (Fig. 2.E, F). This patter is due to the fact that potassium in being a non structural element and is subject to physical removal by leaching (Lousier and Parkinson, 1978; Tisdale *et al.,* 1993; Xu *et al.,* 2004). The concentration of Na was exhibited decreasing trend from the beginning till the end of the study in both forest sites. Sodium releases gradually upto the end of the experiment in both forest sites. Released of Na (99%) in forest site I is higher than site II (96%).

The release of nutrients in forest site I and site II was in the order K > Na > P > N. The release of nutrient elements in the order of K > P > N also reported by Hart *etal.,*1992 in *Ponderosa pine* needle litter at California.

Lignin concentration increases during decomposition preceded in both forest sites. It may be due to by an interaction of factors limiting the activity of the microbial decomposers and possibly a difference between microbial communities. The fungi and bacteria can easily degraded cellulose and hemicellulose than lignin. Lignin is a less energy rich and thus not primarily attacked and degraded more slowly. These would be a proportionally higher decomposition rate of the cellulose and the lignin concentration would increase more quickly. Berg et. al., (1993) and Pandey et. al., (2007) reported increased in lignin concentration in decomposing Scot pine needle litter and Quercus species leaf litter.

Acknowledgements
The Head Department of Botany (P.G.), D.M. College of Science, Imphal, is thankful for provided laboratory facilities and the Conservator, Forest Department, Government of Manipur, Imphal West district is also highly grateful for allowing the study areas.

REFERENCES

1. *Akoto, D. S., Partey, S. T., Abugre, S., Akoto, S., Denich, M., Borgemeister, C. (2022). Comparative analysis of leaf litter decomposition and nutrient release patterns of bamboo and traditional species in agroforestry system in Ghana. Clean. Mater. 4:100068. doi: 10.1016/j.clema.2022.100068*
2. *Anderson, J.M., Ingram J.S.I., 1993. Tropical soil biology and Fertility; a Handbook of Methods. CAB International, Oxford.*
3. *Arunachalam, A., Singh N.D., 2004. Decomposition of Mesua ferrea litter in humid tropics of Arunachal Pradesh , India .Journal of Tropical Forest Science* ***16, 151-159.***
4. *Berg, B., Ekbohm, G., 1991. Litter mass-loss rates and decomposition patterns in some needle and leaf litter types. Long-term decomposition in scots pine forests VII. Canadian Journal of Botany* ***69, 1449 _ 1456.***

ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188

5. *Berg, B., Ekbohm, G., Johansson, M.B., Mc Caugherty, C., Rutigliano, F.A., Virgo de Santo, A., 1996. Maximum decomposition limits of forest litter types; a synthesis. Canadian journal of Botany* ***74, 659 - 672.***
6. *Berg, B., Staaf, H. 1980. Decomposition rate and chemical changes of Scots pine needle litter. II. Influence of chemical composition. Structure and function of Northernconiferous forest - on ecosystem study (Ed. by T. Persson), Ecological Bulletins****32, 373 _ 390.***
7. *Bijayalaxmi, N.D., 2005. Litterfall, litter decomposition and soil microbial biomass dynamics in the mixed oak forest at Langol Hill range, Manipur. Ph.D. Thesis. Manipur University.*
8. *Blair, J.M., 1988. Nitrogen, sulphur and phosphorus dynamics in decomposing deciduousleaf litter in the Southern Appalachians. Soil Biology and Biochemistry****20, 693 _ 701***
9. *Bockheim, J.G., Jepsen, E.A., Heisey, D.M., 1991. Nutrient dynamics in decomposing leaf litter of four tree species on a sandy soil in north western Wisconsin. Canadian Journal of Forest Research* ***21, 803-812.***
10. *Bradford, M.A., Tordof, G.M., Eggers, T., Jones, H. , Newington, J.E. 2002. Microbiota, fauna and mesh size interaction in litter decomposition. Oikos* ***99, 317 _ 323.***
11. *Constantinides, M. , Fownes, J.H., 1994. Nitrogen mineralization from leaves and litter of tropical plants, relationship to nitrogen, lignin and soluble polyphenol concentrations. Soil Biology and Biochemistry* ***26, 49 _ 55.***
12. *Costa AN, Souza JR, Alves KM, et al. (2020) Linking the spatiotemporal variation of litterfall to standing vegetation biomass in brazilian savannas. J Plant Ecol 13:517–524.*
13. *Couteaux, M.M., Bottner, P., Berg, B., 1995. Litter decomposition, climate and litter quality. Trends in ecology and evolution* ***10, 63 _ 66.***
14. *Couteaux, M.M., Mc Tiernan, K., Berg, B., Szuberla, D. , Dardenner, P., 1998. Chemical composition and carbon mineralisation potential of Scots pine needle at different stages of decomposition. Soil Biology and Biochemistry* ***30, 583 _ 595.***
15. *Cusack , D.F.,Chou W.W., Yang W.H., Harmon M.E., Silver W.L., Team, T. L., 2009. Controls on long -term root and leaf litter decomposition in neotropical forest. Global Change Biology* ***15,1339-1355.***
16. *Das, A.K., Ramakrishnan, P.S., 1985. Litter dynamics in Khasi pine of north-east India. Forest Ecology and Management* ***10, 135 _ 154.***
17. *David, R.C., Philip, I., Clives, W., Trevor. G.P. 1998. Decompostion of Pinus sylvestris litter in litter bags: influence of underlying native litter layer. Soil Biology and Biochemistry* ***30, 47 _55.***
18. *Finzi, A.C., Allen, A.S., De Lucia, E.H., Ellsworth, D.S., Schlesinger, W.H., 2001. Forest litter production, chemistry and decomposition following two-years of free airCO_2 enrichment. Ecology* ***82, 470 _ 484.***
19. *García-Palacios P, Mckie BG, Handa IT, et al. (2016a). The importance of litter traits and decomposers for litter decomposition: a comparison of aquatic and terrestrial ecosystems within and across biomes. Funct Ecol 30:819–829.*
20. *Ganjegunte, G.K., Condron, L.M., Clinton, P.W., Davis, M. R., Mahicu, N. 2004. Decomposition and nutrient release from radiata pine (Pinusradiata) coarse woody debris. Forest Ecology and Management* ***187, 1987 _ 211.***
21. *Gilbert, D.J., Bocock,L.K. 1962. Some methods of studying the disappearance anddecomposition of leaf litter. Progress in soil zoology (Ed. by P.W. Murphy). pp* ***348-352.***
22. *Goma-Tchimbakala J., Bernhard - Reversat, F., 2006. Comparison of litter dynamics in three planations of an indigenous timber tree species (Terminalia superba) and anatural tropical forest in Mayombe, Congo. Forest Ecology and Management****229, 304-313.***
23. *Gonzalez, G., Seastedt, T.R., 2001. Soil fauna and plant litter decomposition in tropical and subalpine forests. Ecology* ***82, 955 _ 964.***
24. *Hart, S.C., Firestone, M.K., Paul, E.A., 1992. Decomposition and nutrient dynamics of ponderosa pine needle in a Mediterranean- type climate. Canadian Journal of Forest Research* ***22, 306 _ 314.***
25. *Jia BR, Xu ZZ, Zhou GS, et al. (2018) Statistical characteristics of forest litterfall in China. Sci China Life Sci 61:358–360.*
26. *Jorgenson, R.G., Myer, B., 1990. Nutrient changes in decomposing beechleaflitter assessed using a solution flux approach. Journal of Soil Science* ***41, 279 _ 293.***
27. *Kaushal, R., Verma, K.S. ,Singh K.N., 2006. Decomposition of pattern and nutrients dynamics in leaf litter of Populus deltoides Marsh in Himachal Pradesh. Indian Forester* ***4, 457 _ 466.***
28. *Keyser, P., Kirk T.K., Zeikus J.G., 1978. Lignolytic enzyme of Phanaerochaete chrysosporium synthetized in the absence of lignin in response to nitrogen starvation. Journal of Bacteriology* ***135, 780 _ 797.***
29. *Kocky, M. ,Wilson, S.D., 1997. Litter decomposition and nitrogen dynamics in aspen forest and mixed grass prairie. Ecology* ***78, 732 _ 739.***
30. *Kurz, B. C., Couteaux, M-M., Jean, M.T., Berg, B. , Jean, R. 2005. Acomparison of litter bag and direct observation methods of Scots pineneedle decomposition measurement. Soil Biology and Biochemistry****37, 2315 _ 2318.***
31. *Lousier, J.D., Parkinson, D., 1976. Litter decomposition in a cool temperate deciduous forest. Canadian Journal of Botany* ***54, 419 _ 436.***
32. *Lousier, J.D., Parkinson, D., 1978. Chemical element dynamics in decomposing leaf litter. Canadian Journal of Botany* ***56, 279 - 2812.***
33. *Mc Brayer, J.F.,Cromark, K.Jr.,1980.Effect of snow-pack on Oak-litterbreakdown and nutrient release in a Minnesota forest. Pedobiologia* ***20, 47 _54.***
34. *Marchante, E., Marchante, H., Freitas, H., Kjoller, A., and Struwe, S. (2019). Decomposition of an N-fixing invasive plant compared with a native species: Consequences for ecosystem. Appl. Soil Ecol. 138, 19–31. doi: 10.1016/j.apsoil. 2019.02.016McClaugherty, C.A., Berg, B. 1987. Cellulose, lignin and nitrogen concentrations as rate regulating factors in late stages of forest litter decomposition. Pedobiologia****30,101 _ 112.***
35. *Meentemeyer, V., 1978. Macroclimate and lignin control of litter decomposition rates. Ecology* ***59, 465 _ 472.***
36. *Melillo, J.M., Aber, J.D., Muratore, J.F., 1982. Nitrogen and lignin control of hardwood leaf litter decomposition dynamics Ecology* ***63, 621 _ 626.***
37. *Misra, M.K., Nisanka, S.K. 1997. Litterfall, decomposition and nutrient release in Casuarina equisetifolia plantations on the sandy coast of Orrisa, India. Tropical Ecology* ***38, 109 _ 119.***
38. *Mun, H.T., 2009.Weight loss and Nutrient Dynamics during leaf litter decomposition of Quercus mongolica in*

ISSN (Online): 2455-3662
EPRA International Journal of Multidisciplinary Research (IJMR) - Peer Reviewed Journal
Volume: 12| Issue: 1| January2026|| Journal DOI: 10.36713/epra2013 || SJIF Impact Factor 2025: 8.691 || ISI Value: 1.188

Mt. Worak National Park. Journal of Ecology and Field Biology ***32(2), 123-127.***

39. *Mtambanengwe, F., Kirchmann, H., 1995. Litter from a tropical savanna woodland (miombo) : chemical composition and C and N mineralization. Soil Biology and Biochemistry* ***27, 1639 _ 1651.***
40. *Namgung, J., Han A. R., Mun, H. T., 2008. Weight loss and nutrient dynamics during leaf litter decomposition of Quercus variabilis and Pinus densiflora at Mt. Worak National Park . Journal of Ecology and Field Biology* ***31, 291 -295.***
41. *Olson, J.S., 1963. Energy storage and balance of producers and decomposers in ecological systems Ecology* ***44, 322 _ 331.***
42. *Palm, C.A., Sanchez, P.A., 1991. Nitrogen release from the leaves of some tropical legumes as affected by their lignin and polyphenolic contents. Soil Biology and Biochemistry* ***23, 83 _ 88.***
43. *Pandey, R.R., Sharma, G., Tripathi, S.K., Singh, A.K. 2007. Litterfall, litter decoposition and nutrient dynamics in a subtropical oak forest and managed plantation in northeastern India. Forest Ecology and Management* ***240, 96-104.***
44. *Peach, K., Tracey, M.V., 1955. Modern methods of plant analysis.Vol.I:springer, Berlin.*
45. *Sarjubala, A.D., 2002. Litter decomposition and nutrient mineralization in the dry tropical deciduous forest ecosystem of Manipur. Ph.D. Thesis. Manipur University.*
46. *Sharma, A.P., Bidhan, R.P., Bisht, J.S., 1984. Microarthropods in litter bags p. 299-304. In: J.S. Singh and S.P. Singh (eds.) Integrated ecological study of eastern Kumaun Himalaya with emphasis on national resources. Department of Science and Technology, New Delhi.*
47. *Tisdale. Smuel. L., Nelson. W. L., Benaton. J. D., Havlin. J. L.,1993. Soil Fertility and Fertilisers. fifth ed Prentice Hall of India Private Limited. New Delhi pp.* ***53-60.***
48. *Tripathi, S.K., Singh, K.P., 1992. Nutrient immobilization and release patterns during plant decomposition in a dry tropical bamboo savana, India. Biology and Fertility of Soils* ***14, 191 _ 199.***
49. *Upadhyay, V.P., Singh, J.S., 1989. Patterns of nutrient immobilization and releaseindecomposing forest litter in Central Himalaya, India. Journal of Ecology****77, 127 _ 146.***
50. *Upadhyay, V.P., Singh, J.S., Meentemeryer, V., 1989. Dynamics and weight loss of leaf litter in Central Himalayan forests: Abiotic versus litter quality influences. Journal of ecology* ***77, 147 _ 161.***
51. *Up-degraff's, D.M., 1969. Semi-micro determination of cellulose in biological materials. Annals of Biochemistry* ***32, 430 _ 434.***
52. *Wardle, D.L., Nelson, M., Zackrisson, O., Gallel, C., 2003. Determinants of litter mixing effects in a Swedish boreal forest. Soil Biology and Biochemistry 35, 827 _ 835.*
53. *Webderburn, M.E., Carter, J., 1999. Litter decomposition by four tree types for use in silvipastoral systems. Soil Biology and Biochemistry* ***31, 455 _ 461.***
54. *Xu, X., Hirata, E., Shibata, H., 2004. Effect of Typhoon disturbance on fine litterfall and related nutrient input in a subtropical forest on Okinawa Island, Japan, Basic and Applied Ecology* ***5, 271 _ 282.***