



ON THE HOMOGENEOUS TERNARY QUADRATIC EQUATION

$$7x^2 + 3y^2 = 220z^2$$

¹**S .Mallika**

¹Assistant professor,
 Department of Mathematics,
 Shrimati Indira Gandhi College,
 Trichy, Tamil Nadu,
 India

²**G.Annes joshiba**

²P G Scholar,
 Department of Mathematics,
 Shrimati Indira Gandhi College,
 Trichy, Tamil Nadu,
 India

ABSTRACT

The ternary quadratic equation given by $7x^2 + 3y^2 = 220z^2$ is considered and searched for its many different integer solution. Five different choices of integer solution of the above equations are presented. A few interesting relations between the solutions and special polygonal numbers are presented.

KEY WORDS: ternary quadratic, integer solutions

MSC subject classification: 11D09

1. INTRODUCTION

The Diophantine equations offer an unlimited field for research due to their variety [1-3]. In particular, one may refer [4-8] for quadratic equations with three unknowns. This communication concerns with yet another interesting equation $7x^2 + 3y^2 = 220z^2$ representing homogeneous equation with three for determining its infinitely many non-zero integral points. Also, few interesting relations among the solutions are presented.

2. NOTATIONS

- $t_{m,n} = n^{th}$ term of a regular polygon with m sides.

$$= n \left(1 + \frac{(n-1)(m-2)}{2} \right)$$

- $p_{rn} =$ pronic number of rank n



$$= n(n + 1)$$

3 .METHOD OF ANALYSIS:

The Quadratic Diophantine equation with three unknowns to be solved is given by

$$7x^2 + 3y^2 = 220z^2 \tag{1}$$

Consider the linear transformation

$$\left. \begin{aligned} x &= X - 3T \\ y &= X + 7T \end{aligned} \right\} \tag{2}$$

Substituting (2) in (1) we get,

$$7(X^2 - 9T^2) + 3(X^2 + 49T^2) = 220T^2$$

$$7X^2 + 63T^2 + 3X^2 + 147T^2 = 220T^2$$

$$10X^2 + 210T^2 = 220T^2$$

$$X^2 + 21T^2 = 22T^2 \tag{3}$$

(3) Is solved through different approaches and the different patterns of solutions of (1) obtained are presented below.

PATTERN: 1

Assume

$$z = a^2 + 21b^2$$

(3) Can also be written as

$$X^2 + 21T^2 = 22 * T^2 \tag{4}$$

Write '22' as

$$22 = (1 + i\sqrt{21})(1 - i\sqrt{21})$$



In equation (4), we get

$$(X + i\sqrt{21}T)(X - i\sqrt{21}T) = (a + i21b)^2(a - i\sqrt{21}b)^2(1 + i\sqrt{21})(1 - i\sqrt{21})$$

Consider the positive factor

$$X + i\sqrt{21}T = (a + i\sqrt{21}b)^2(1 + i\sqrt{21})$$

$$X + i\sqrt{21}T = (a^2 - 21b^2 + i2ab\sqrt{21})(1 + i\sqrt{21})$$

$$X + i\sqrt{21}T = (a^2 - 21b^2 - 42ab) + i\sqrt{21}(a^2 - 21b^2 + 2ab)$$

Equating real and imaginary parts

$$\left. \begin{aligned} X &= a^2 - 21b^2 - 42ab \\ T &= a^2 - 21b^2 + 2ab \end{aligned} \right\} \quad (5)$$

From (2), we obtain the non-zero distinct integral solution of (1) as

$$x(A, B) = -2a^2 + 42b^2 - 48ab$$

$$y(A, B) = 8a^2 - 168b^2 - 28ab$$

$$z(A, B) = a^2 + 21b^2$$

PROPERTIES:

$$1. x(A,1) + y(A,1) - 82t_{4,a} + 76p_{ra} \equiv 0 \pmod{2}$$

$$2. y(A,1) + z(A,1) - 37t_{4,a} + 28p_{ra} \equiv 0 \pmod{3}$$

$$3. x(A,1) - y(A,1) - 10t_{4,a} + 20p_{ra} = 210$$

PATTERN: 2

In (4) '22' can also written as

$$22 = \frac{(19 + 3i\sqrt{21})(19 - 3i\sqrt{21})}{5^2} \quad (6)$$



Substituting (4) & (6) in (3) and employing the method of factorization, we get

$$(X + i\sqrt{21}T)(X - i\sqrt{21}T) = (a + i\sqrt{21}b)^2 (a - i\sqrt{21}b)^2 \frac{(19 + 3i\sqrt{21})(19 - 3i\sqrt{21})}{5^2}$$

Consider the positive factor

$$X + i\sqrt{21}T = \frac{19 + 3i\sqrt{21}}{5} (a + i\sqrt{21}b)^2$$

$$X + i\sqrt{21}T = \frac{19 + 3i\sqrt{21}}{5} (a^2 - 21b^2 + i2\sqrt{21}ab)$$

$$X + i\sqrt{21}T = (19a^2 - 399b^2 - 126ab) + i\sqrt{21}(3a^2 - 63b^2 + 38ab)$$

Equating real and imaginary parts of the above equation, we get

$$X = \frac{19a^2 - 399b^2 - 126ab}{5}$$

$$Y = \frac{3a^2 - 63b^2 + 38ab}{5}$$

From (2)

$$x = 2a^2 - 42b^2 - 48ab$$

$$y = 8a^2 - 168b^2 + 28ab$$

We obtain the non-zero distinct integral solution of (1) as

$$x(A, B) = 2a^2 - 42b^2 - 48ab$$

$$y(A, B) = 8a^2 - 168b^2 + 28ab$$

$$z(A, B) = a^2 + 21b^2$$



PROPERTIES:

1. $x(A,1) + y(A,1) - 30t_{4,a} + 20p_{ra} + 210 = 0$
2. $x(A,1) + z(A,1) - 51t_{4,a} + 48p_{ra} \equiv 0 \pmod{7}$
3. $y(A,1) + z(A,1) + 19t_{4,a} - 28p_{ra} + 147 = 0$

PATTERN: 3

Write the equation (4) as

$$X^2 + 21T^2 = 22z^2 * 1 \tag{7}$$

'1' can also be written as

$$1 = \frac{(2 + i\sqrt{21})(2 - i\sqrt{21})}{5^2} \tag{8}$$

Write '22' as

$$22 = (1 + i\sqrt{21})(1 - i\sqrt{21}) \tag{9}$$

Substituting (7), (8) & (9) in (3) and employing the method of factorization, we get

$$(X + i\sqrt{21}T)(X - i\sqrt{21}T) = \left[(a + i\sqrt{21}b)^2 (a - i\sqrt{21}b)^2 (1 + i\sqrt{21})(1 - i\sqrt{21}) \right] * \left[\frac{(2 + i\sqrt{21})(2 - i\sqrt{21})}{5^2} \right] \text{ Consider}$$

r the positive factor

$$(X + i\sqrt{21}T) = \left[(a + i\sqrt{21}b)^2 (1 + i\sqrt{21}) \right] * \left[\frac{(2 + i\sqrt{21})}{5} \right]$$

$$(X + i\sqrt{21}T) = \left[(a^2 - 21b^2 + i2ab\sqrt{21})(1 + i\sqrt{21}) \right] * \left[\frac{(2 + i\sqrt{21})}{5} \right]$$

$$(X + i\sqrt{21}T) = \frac{(-19a^2 + 399b^2 - 126ab) + i\sqrt{6}(3a^2 - 63b^2 - 38ab)}{5}$$



Equating real and imaginary parts of the above equation, we get

$$X = \frac{-19a^2 + 399b^2 - 126ab}{5}$$

$$T = \frac{3a^2 - 63b^2 - 38ab}{5}$$

From (2)

$$x = \frac{-28a^2 + 588b^2 - 12ab}{5}$$

$$y = \frac{2a^2 - 42b^2 - 392ab}{5}$$

Assume $a=5A$, $b=5B$ in the above equations, we obtain the non-zero integral solution of (1) as

$$x(A, B) = -140A^2 + 2940B^2 - 60AB$$

$$y(A, B) = 10A^2 - 210B^2 - 1960AB$$

$$z(A, B) = 25A^2 + 525B^2$$

PROPERTIES:

1. $x(A,1) + y(A,1) - 1890t_{4,A} + 2020p_{ra} \equiv 0 \pmod{5}$
2. $x(A,1) + z(A,1) + 55t_{4,A} + 60p_{ra} = 3465$
3. $y(A,1) + z(A,1) - 1995t_{4,A} + 1960p_{ra} \equiv 0 \pmod{5}$

PATTERN: 4

'1' can also written as

$$1 = \frac{(2 + 5i\sqrt{21})(2 - 5i\sqrt{21})}{23^2} \tag{10}$$



Substituting (10) and (9) in (7) and employing the method of factorization, we get

$$(X + i\sqrt{21}T)(X - i\sqrt{21}T) = \left[(a + i\sqrt{21}b)^2 (a - i\sqrt{21}b)^2 (1 + i\sqrt{21})(1 - i\sqrt{21}) \right] * \frac{(2 + 5i\sqrt{21})(2 - 5i\sqrt{21})}{23^2} \text{ Consider}$$

the positive factor

$$(X + i\sqrt{21}T) = \left[(a^2 - 21b^2 + i2\sqrt{21}ab)(1 + i\sqrt{21}) \right] * \left[\frac{(2 + 5i\sqrt{21})}{23} \right]$$

$$(X + i\sqrt{21}T) = \left[\frac{(7i\sqrt{21} - 103)(a^2 - 21b^2 + i2\sqrt{21}ab)}{23} \right]$$

$$(X + i\sqrt{21}T) = \frac{(-103a^2 + 2163b^2 - 294ab) + i\sqrt{21}(7a^2 - 147b^2 - 206ab)}{23}$$

Equating real and imaginary parts of the above equation, we get

$$X = \frac{-103a^2 + 2163b^2 - 294ab}{23}$$

$$T = \frac{7a^2 - 147b^2 - 206ab}{23}$$

From (1) we get

$$x = \frac{-124a^2 + 2604b^2 + 324ab}{23}$$

$$y = \frac{-103a^2 + 2163b^2 - 294ab}{23}$$

Assume a=23A, b=23B in the above equation, we obtain the non-zero distinct integral solution (1) as

$$x(A, B) = -2852A^2 + 59892B^2 + 7452AB$$

$$y(A, B) = -1242A^2 + 26082B^2 - 39928AB$$



$$z(A, B) = 529A^2 + 11109B^2$$

PROPERTUES:

1. $x(A,1) + z(A,1) + 9775t_{4,A} - 7452p_{rA} = 71001$
2. $x(A,1) - z(A,1) + 10833t_{4,A} - 7452p_{rA} \equiv 0 \pmod{3}$
3. $y(A,1) + z(A,1) - 39215t_{4,A} + 39928p_{rA} = 37191$

PATTERN: 5

(3) Can also written as

$$\begin{aligned} X^2 - z^2 &= 21(z^2 - T^2) \\ (X + z)(X - z) &= 21(z + T)(z - T) \end{aligned} \tag{11}$$

Case: 1

(3) Can be written as in the form of ratio as

$$\frac{X + z}{z + 1} = \frac{21(z - T)}{X - z} = \frac{\alpha}{\beta} \tag{12}$$

which is equivalent to the system of double equations as

$$\left. \begin{aligned} \beta X - \alpha T + (-\alpha + \beta)z &= 0 \\ -\alpha X - 21\beta T + (\alpha + 21\beta)z &= 0 \end{aligned} \right\} \tag{13}$$

Solving (13) by method of cross multiplication, we get

$$\left. \begin{aligned} X &= -\alpha^2 + 21\beta^2 - 42\alpha\beta \\ T &= \alpha^2 - 21\beta^2 - 2\alpha\beta \\ z &= -\alpha^2 - 21\beta^2 \end{aligned} \right\} \tag{14}$$

Substituting (14) in (2) and the non-zero distinct integer solution of (1) are given by

$$x(\alpha, \beta) = -4\alpha^2 + 84\beta^2 - 36\alpha\beta$$



$$y(\alpha, \beta) = 6\alpha^2 - 126\beta^2 + 56\alpha\beta$$

$$z(\alpha, \beta) = -\alpha^2 - 21\beta^2$$

PROPERTIES:

1. $x(\alpha, 1) + y(\alpha, 1) - 94t_{4,\alpha} + 92p_{r\alpha} + 42 = 0$
2. $x(\alpha, 1) + z(\alpha, 1) - 31t_{4,\alpha} + 36p_{r\alpha} \equiv 0 \pmod{3}$
3. $y(\alpha, 1) + z(\alpha, 1) - 61t_{4,\alpha} + 56p_{r\alpha} + 147 = 0$

Case: 2

Equation (11) can be written as

$$\frac{X + z}{21(z - T)} = \frac{z + T}{X - z} = \frac{\alpha}{\beta} \tag{15}$$

Which is equivalent to the system of double equation as

$$\left. \begin{aligned} \beta X + 21\alpha T + (-21\alpha + \beta)z &= 0 \\ -\alpha X + \beta T + (\alpha + \beta)z &= 0 \end{aligned} \right\} \tag{16}$$

Solving (16) by method cross multiplication, we get

$$\left. \begin{aligned} X &= 21\alpha^2 - \beta^2 + 42\alpha\beta \\ T &= 21\alpha^2 - \beta^2 - 2\alpha\beta \\ z &= 21\alpha^2 + \beta^2 \end{aligned} \right\} \tag{17}$$

Substituting (17) in (2), the non-zero distinct integral solution of (1) is given by

$$x(\alpha, \beta) = -42\alpha^2 + 2\beta^2 + 48\alpha\beta$$

$$y(\alpha, \beta) = 168\alpha^2 - 8\beta^2 + 28\alpha\beta$$

$$z(\alpha, \beta) = 21\alpha^2 + \beta^2$$



PROPERTIES:

1. $x(\alpha,1) + y(\alpha,1) - 50t_{4,\alpha} - 76p_{r\alpha} + 6 = 0$
2. $x(\alpha,1) + z(\alpha,1) + 69t_{4,\alpha} - 48p_{r\alpha} \equiv 0 \pmod{3}$
3. $y(\alpha,1) + z(\alpha,1) - 161t_{4,\alpha} - 28p_{r\alpha} + 7 = 0$

Case: 3

(11) can also be written in the form of the ratio as

$$\frac{X + z}{7(z + T)} = \frac{3(z - T)}{X - z} = \frac{\alpha}{\beta} \quad (18)$$

which is equivalent to the system of double equations as

$$\left. \begin{aligned} \beta X - 7\alpha T + (-7\alpha + \beta)z &= 0 \\ -\alpha X - 3\beta T + (\alpha + 3\beta)z &= 0 \end{aligned} \right\} \quad (19)$$

Solving (19) by method of cross multiplication, we get

$$\left. \begin{aligned} X &= -7\alpha^2 + 3\beta^2 - 42\alpha\beta \\ T &= 7\alpha^2 - 3\beta^2 - 2\alpha\beta \\ z &= -7\alpha^2 - 3\beta^2 \end{aligned} \right\} \quad (20)$$

Substituting (20) in (2), the non-zero distinct integral solution of (1) is given by

$$\begin{aligned} x(\alpha, \beta) &= -28\alpha^2 + 12\beta^2 - 36\alpha\beta \\ y(\alpha, \beta) &= 42\alpha^2 - 18\beta^2 - 56\alpha\beta \\ z(\alpha, \beta) &= -7\alpha^2 - 3\beta^2 \end{aligned}$$

PROPERTIES:

1. $x(\alpha,1) + y(\alpha,1) - 106t_{4,\alpha} + 92p_{r\alpha} + 6 = 0$



$$2. x(\alpha,1) + z(\alpha,1) - t_{4,\alpha} + 36p_{r\alpha} \equiv 0 \pmod{9}$$

$$3. y(\alpha,1) + z(\alpha,1) - 91t_{4,\alpha} + 56p_{r\alpha} + 21 = 0$$

Case: 4

(11) can be written in the form of the ratio as

$$\frac{X + z}{3(z + T)} = \frac{7(z - T)}{X - z} = \frac{\alpha}{\beta} \quad (21)$$

which is equivalent to the system of double equations is

$$\left. \begin{aligned} \beta x - 3\alpha T + (-3\alpha + \beta)z &= 0 \\ -\alpha X - 7\beta T + (\alpha + 7\beta)z &= 0 \end{aligned} \right\} \quad (22)$$

solving (22) by method of cross multiplication, we get

$$\left. \begin{aligned} X &= 3\alpha^2 - 7\beta^2 + 42\alpha\beta \\ u &= -3\alpha^2 + 7\beta^2 + 2\alpha\beta \\ T &= 3\alpha^2 + 7\beta^2 \end{aligned} \right\} \quad (23)$$

Substituting (23) in (2), the non-zero distinct integral solution of (1) is given by

$$x(\alpha, \beta) = 12\alpha^2 - 28\beta^2 + 36\alpha\beta$$

$$y(\alpha, \beta) = -18\alpha^2 + 42\beta^2 + 56\alpha\beta$$

$$z(\alpha, \beta) = 3\alpha^2 + 7\beta^2$$

PROPERTIES:

$$1. x(\alpha,1) + y(\alpha,1) + 98t_{4,\alpha} - 92p_{r\alpha} \equiv 0 \pmod{7}$$

$$2. x(\alpha,1) + z(\alpha,1) + 21t_{4,\alpha} - 36p_{r\alpha} + 21 = 0$$

$$3. y(\alpha,1) + z(\alpha,1) + 71t_{4,\alpha} - 56p_{r,\alpha} = 49$$



Case: 5

(11) can be written in the form of the ratio as

$$\frac{X - z}{7(z - T)} = \frac{3(z + T)}{X + T} = \frac{\alpha}{\beta} \quad (24)$$

which is equivalent to the system of double equations as

$$\left. \begin{aligned} \beta X + 7\alpha T - (7\alpha + \beta) &= 0 \\ -\alpha X + 3\beta T + (-\alpha + 3\beta) &= 0 \end{aligned} \right\} \quad (25)$$

Solving (25) by method of cross multiplication, we get

$$\left. \begin{aligned} X &= -7\alpha^2 + 3\beta^2 + 42\alpha\beta \\ T &= 7\alpha^2 - 3\beta^2 + 2\alpha\beta \\ z &= 7\alpha^2 + 3\beta^2 \end{aligned} \right\} \quad (26)$$

Substituting (26) in (2), the non-zero distinct integral solutions of (1) are given by

$$x(\alpha, \beta) = -28\alpha^2 + 12\beta^2 + 36\alpha\beta$$

$$y(\alpha, \beta) = 42\alpha^2 - 18\beta^2 + 56\alpha\beta$$

$$z(\alpha, \beta) = 7\alpha^2 + 3\beta^2$$

PROPERTIES:

$$1. x(\alpha, 1) + y(\alpha, 1) + 78t_{4,\alpha} - 92p_{r\alpha} + 6 = 0$$

$$2. x(\alpha, 1) + z(\alpha, 1) + 57t_{4,\alpha} - 36p_{r\alpha} \equiv 0 \pmod{5}$$

$$3. y(\alpha, 1) + z(\alpha, 1) + 7t_{4,\alpha} - 56p_{r\alpha} + 15 = 0$$

CONCLUSION

In this paper, we have made an attempt to obtain infinitely many non-zero distinct integer solutions to special ternary quadratic Diophantine equation $7x^2 + 3y^2 = 220z^2$. In each case, a few interesting relations among the solution are presented. As the Diophantine equations are



rich in variety, one may search for integer solutions to higher degree Diophantine equations with multiple variables along with suitable properties.

REFERENCES

1. R.D Carmichael, *Theory of Numbers and Diophantine Analysis*, Dover Publications New York, 1950.
2. L.E.Dickson, *History of theory of numbers, Vol. II Chelsea Publishing Co., New York, 1952*
3. L.J.Mordell, *Diophantine Equations Academic Press, London, 1969*
4. Dr.S.Mallika, D.Maheshwari, R.Anbarasi on the homogeneous Ternary quadratic Diophantine equation $16x^2+18xy+3y^2=11z$. *Infokara research journal volume 9, issue 4, April 2020 Page 26-31.*
5. Dr.S.Mallika, Dr.V.Praba observation on the homogeneous ternary quadratic Diophantine equation $25x^2-20xy+10y^2=7z^2$ *journal of interdisciplinary cycle research volume 12, issue 4, April 2020 page 1301-1308*
6. Dr.s.Mallika,T.Arthi “on the Homogeneous Ternary quadratic Diophantine equation $9x^2 -6xy+2y^2 =14z^2$ ” (IJRAR)*International Journal of Research and Analytical Review, Volume 7,issue 1 page 937-945 March 2020*
7. D.Hema, S.Mallika, on a ternary quadratic Diophantine equation $5y^2=3x^2+2z^2$ ” *Journal of Mathematics and Informatics, volume 10, 2017, 157-165.*
8. Dr.S.Mallika, “On the homogeneous ternary quadratic Diophantine equation $6x^2 + 7y^2 = 559z^2$ ” ,*International Journal of mathematics trends and technology volume 65, issue 7, July 2019 ,page 206-217.*