SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

PROPERTIES OF τ AND σ FUNCTIONS

Hamant Kumar[1], Vishnu Kant[2]
[1]*Department of Mathematics*
[2]*Department of English*
V. A. Govt. Degree College, Atrauli, Aligarh, Uttar Pradesh (India)

ABSTRACT
In this article, we study τ and σ functions and their properties. Utilizing τ and σ functions, we investigate perfect number, deficient numbers, abundant numbers, amicable numbers, friendly numbers and solitary numbers.

1. INTRODUCTION

τ (tau) and σ (sigma) functions are most important number theoretic functions also known as arithmetic functions. Arithmetic functions are defined for all positive integers. τ and σ functions belong to a large class of arithmetic functions called multiplicative functions and study of some their fascinating properties. τ function, also known as the divisor function, determines the number of positive divisors of a given integer and the σ function also known as the sum of divisors function, calculates the sum of all positive divisors of a given integer. These functions play a significant role in various areas of number theory including the study of prime factorization, divisor properties and properties of perfect numbers. We can use the σ function to study a marvelous class of numbers called perfect numbers the term perfect numbers. The term perfect numbers was coined by Pythagoreans. Euclid shown that every integer $N = 2^{n-1}(2^n - 1)$, where $2^n - 1$ is a prime, is a perfect. Amicable numbers were known to the Pythagoreans, who credited them with many mystical properties.

2. THE FUNCTION $\tau(N)$

2.1 Definition. Let n be a given positive integer. Then $\tau(n)$ is the number of positive divisors of n including the divisors 1 and n.
For an example of these notion, consider n = 12.

The divisors of 12 are 1, 2, 3, 4, 6 and 12.

Hence $\tau(n) = 6$.

For the first few integers,

$\tau(1) = 1$, $\tau(2) = 2$, $\tau(3) = 2$, $\tau(4) = 3$, ……

It is not difficult to see that $\tau(n) = 2$ if and only if n is a prime number.

2.2 Theorem. Let $n = p_1^{a_1}. p_2^{a_2}. p_3^{a_3} \ldots\ldots\ldots p_k^{a_k}$. Then the divisors of n are the various terms of the expansion of

$n = (1 + p_1 + p_1^2 + \ldots\ldots\ldots + p_1^{a_1})\ (1 + p_2 + \ldots\ldots + p_2^{a_2}) \ldots\ldots\ldots (1 + p_k + \ldots\ldots + p_k^{a_k})$.

2.3 Theorem. If $n = p_1^{k_1} p_2^{k_2} \ldots\ldots\ldots\ldots P_r^{k_r}$ is the prime factorization of $n > 1$, then

(1) $\tau(n) = (k_1 + 1)(k_2 + 1) \ldots\ldots\ldots\ldots\ldots (k_n + 1)$ and
(2) $\sigma(n) = \frac{p1^{k1+1}-1}{p1-1}\ \frac{p2^{k2+1}-1}{p2-1} \ldots\ldots\ldots\ldots\ldots \frac{pr^{kr+1}-1}{pr-1}$.

2.4 Example. Let $n = 2700 = 2^2 \times 3^3 \times 5^2$. Then

SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

$\tau(2700) = (2 + 1)\ (3 + 1)\ (2 + 1) = 36$

$\sigma(2700) = \frac{2^3-1}{2-1} \times \frac{3^4-1}{3-1} \times \frac{5^3-1}{5-1} = 8680$

2.5 Theorem. τ(n) is odd if and only if n is a perfect square.

Proof. Suppose $n = p_1^{a_1}.\ p_2^{a_2}.\ p_3^{a_3} \ldots\ldots\ p_k^{a_k}$.

(1) Let n be square integer. Then $a_1, a_2, \ldots\ldots a_k$ are all even. Hence

$\tau(n) = (a_1 + 1)\ (a_2 + 1) \ldots\ldots\ldots\ldots\ldots (a_k + 1)$
= product of odd factors omly
= an odd integer.
(2) Let τ(n) be odd. Then $(a_1 + 1)\ (a_2 + 1) \ldots\ldots\ldots\ldots\ldots (a_k + 1)$ is odd. This implies that $a_1, a_2, \ldots . a_k$ are all even. It follows that n is a square integer.

2.6 Corollary. Let τ(n) is even if and only if n is not a square integer.

2.7 Definition. A number theoretic function f is said to be multiplicative if

$$f(m\ n) = f(m)\ f(n)$$

whenever, gcd(m, n) = 1.

The function τ and σ are both multiplicative functions.

2.8 Definition. The divisors of an integer n, excluding the divisor n are called proper divisors of n, and the sum of all proper divisors of n is denoted by $\sigma_0(n)$.

Examples

(1) Let n = 12. Then the divisors of 12 are 1, 2, 3, 4, 6, 12. Proper divisors of 12 are 1, 2, 3, 4, 6. Therefore

$\sigma(12) = 1 + 2 + 3 + 4 + 6 + 12 = 28$

$\sigma_0(12) = 1 + 2 + 3 + 4 + 6 = 16.$

(2) Let n = 28. Then

$\sigma(28) = 1 + 2 + 4 + 7 + 14 + 28 = 56$

$\sigma_0(28) = 1 + 2 + 4 + 7 + 14 = 28.$

Here $\sigma(n) = 2n$ and $\sigma_0(n) = n$

2.9 Definition. An integer n is said to be a perfect number if the sum of its proper divisors is equal to n. Thus n is a perfect number of $\sigma_0(n) = n$, or $\sigma(n) = 2n$.

2.10 Example. Let $n = 496 = 2^4 \times 31$. Then

$\sigma(496) = \sigma(2^4)\ \sigma(31) = \frac{2^5-1}{2-1} \times (31 + 1)$

SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

$= 2 \times 496$

So, 496 is a perfect number.

There is a simple theorem which enables us to find even perfect numbers. The theorem bears the name of Euclid.

2.11 Theorem (Euclid). Let $2^k - 1$ be a prime. Then $2^{k-1}(2^k - 1)$ is a perfect number.

Proof. Let $n = 2^{k-1}(2^k - 1)$. Then, since 2^{k-1} is a prime,

$$\sigma(n) = \sigma(2^{k-1}(2^k - 1) = \frac{2^k - 1}{2 - 1} \times (2^k - 1 + 1)$$

$$= (2^k - 1)\, 2^k = 2\left[(2^{k-1}(2^k - 1))\right]$$

Which proves the theorem.

2.12 Definition. n is called a deficient number if $\sigma_0(n) < n$.

Examples

(1) $\sigma(15) = \sigma(5)\,\sigma(3) = 24$. Hence $\sigma_0(15) = 24 - 15 = 9 < 15$.

Therefore 15 is a deficient number.

(2) $\sigma(32) = \sigma(2^5) = 2^6 - 1 = 63$.

Hence $\sigma_0(32) = 63 - 32 = 31 < 32$.

Therefore, 32 is a deficient number.

(3) If p is a prime, there is only one proper devisor of p namely 1. Hence $\sigma_0(p) = 1 < p$. Thus every prime is deficient.

(4) p^k is deficient, for

$$\sigma 0(p^k) = \frac{p^{k-1} - 1}{p - 1} - p^k = \frac{p^k - 1}{p - 1} < p^k$$

(5) If find p and q are primes then pq is deficient.

For, $\sigma 0(pq) = (p + 1)(q + 1) - pq$

$= p + q + 1$

$= pq - [(p - 1)(q - 1) - 2]$

$< pq.$

2.13 Definition. n is called an abundant number if $\sigma_0(n) > n$.

Examples. $\sigma 0(12) = 1 + 2 + 3 + 4 + 6 = 16 > 12.$

Hence 12 is an abundant number.

SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

$$\sigma 0(30) = \sigma(2 \times 3 \times 5) - 30$$

$$= (2+1)(3+1)(5+1) - 30$$

$$= 42 > 30$$

(1) $\sigma(945) = \sigma(3^3 \times 5 \times 7) - 945$

$$= 40 \times 6 \times 8 - 945$$

$$= 975 > 945.$$

Hence 945 is an abundant number. It is worth nothing here that 945 is the first odd abundant number.

2.14 Definition. An integer n is called a multiply perfect number of class k if $\sigma_0(n) = kn$.

Example.

I $\sigma 0(120) = \sigma(2^3 \times 3 \times 5) - 120 = 2 \times 120$

Hence 120 is a multiply perfect number of class 2.

II $\sigma 0(2^9 \times 3 \times 11 \times 31) = (2^{10} - 1) \times 4 \times 12 \times 32 - 2^9 \times 3 \times 11 \times 31$

$$= 2(2^9 \times 3 \times 11 \times 31)$$

Hence $2^9 \times 3 \times 11 \times 31$ is a multiply perfect number of class 2.

2.15 Definition. Let there be two integers a and b such that $\sigma 0(a) = b \; and \; \sigma 0(b) = a$. Then (a, b) is called a pair of amicable numbers.

Example.

I. $\sigma 0(220) = 284 \; and$
$\sigma 0(284) = 220$

Hence (220, 284) is a pair of amicable numbers. This was the only pair discovered before fermate applied his powerful mind to this subject.

II. It was Euler, however, who undertook a systematic study of amicable number and he gave a list of 60 Such pairs some of these are
   a) $(2 \times 5 \times 7 \times 19 \times 107, \quad 2 \times 5 \times 47 \times 350)$
   b) $(2^3 \times 5 \times 251, \quad 2^2 \times 13 \times 107$
   c) $(2^3 \times 17 \times 79, \quad 2^3 \times 23 \times 59)$

2.16 Theorem. A pair of integers (a, b) is amicable if and only if

$$\sigma(a) = \sigma(b) = a + b$$

SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

Proof.

i. Let $\sigma(a) = \sigma(b) = a + b$. Then
$\sigma(a) - a = b \; and \; \sigma(b) - b = a$

This implies that $\sigma 0(a) = b$ and $\sigma 0(b) = a$

Therefore, (a, b) is an amicable pair.

ii. Let (a, b) be an amicable pair. Then
$\sigma 0(a) = b$ and $\sigma 0(b) = a$. Hence
$\sigma(a) = a + b$ and $\sigma(b) = a + b$

This proves the theorem completely.

2.17 Theorem. Let

$a_k = 2^k(3\times 2^{k-1} - 1)\,(3 \times 2^k - 1)$ and

$b_k = 2^k(9 \times 2^{2k-1} - 1)$

Where $k > 1$. Then (a_k, b_k) is an amicable pair provided the integers.

$$3 \times 2^{k-1} - 1, \qquad 3 \times 2^k - 1, \qquad 9 \times 2^{2k-1} - 1 \; are \; primes.$$

Proof. We have

$$a_k + b_k = 2^k(3 \times 2^{k-1} - 1)(3 \times 2^k - 1) + 2^k(9 \times 2^{2k-1} - 1)$$

$$= 2^k(9 \times 2^{2k-1} - 3 \times 2^k - 3 \times 2^{k-1} + 1 + 9 \times 2^{2k-1} - 1)$$

$$= 9 \times 2^{3k} - 6 \times 2^{2k-1} - 3 \times 2^{2k-1}$$

$$= 9 \times 2^{2k-1}(2^{k+1} - 1).$$

$\sigma_0(n) = \sigma(2^k)\;\; \sigma(3 \times 2^{k-1} - 1)\; \sigma(3 \times 2^k - 1)$

Since

$$3 \times 2^{k-1} - 1 \text{ is odd for k1,}$$

$$= (2^{k+1} - 1)(3 \times 2^{k-1})(3 \times 2^k)$$

$$= 9 \times 2^{2k-1}\left(2^{k=1} - 1\right)$$

$= a_k + b_k$

$\sigma(bk) = \sigma(2^k)\; \sigma(9 \times 2^{2k-1} - 1)$

SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

$= (2^{k+1} - 1)\, 9 \times 2^{2k-1}$

$= a_k + b_k$

It follows that (a_k, b_k) is an amicable pair.

There exist only three amicable pairs conforming to the above theorem for $k \leq 200$. These are

S. No.	k	a_k	b_k
1.	2	220	284
2.	4	17296	18416
3.	7	9363584	9437056

We give below a few more examples of amicable pairs. These are

a) (1184, 1210)
b) (2620, 2924)
c) (5020, 5564)
d) (6232, 6368)

2.18 Abundancy Index. Abundancy index is the ratio of a number's sum of divisors to the number itself.

2.19 Friendly Numbers. Friendly numbers are two or more numbers that have the same abundancy index.

Example

i. 6 and 28 are friendly number.

The devisors of

6 are 1, 2, 3, 6 and

28 are 1, 2, 4, 8, 14, 28.

The abundancy index of 6 is $\frac{\sigma(n)}{n} = \frac{1+2+3+6}{6} = \frac{12}{6}$

$=2$

The abundancy index of 28 is $\frac{\sigma(n)}{n} = \frac{1+2+4+7+14+28}{28} = \frac{56}{28} = 2$

Hence the number 6 and 28 are friendly numbers because they both have an abundancy of 2.

SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

ii. 30 and 140 are friendly numbers.
iii. 80 and 200 are friendly numbers.

2.20 Solitary Numbers

The numbers that are not part of any friendly pair are called solitary numbers.

Note: Every prime is a solitary number.

Be have our of $\tau(n)$ as n $\to \infty$

I. We know that ever integer $n > 1$ has at least two divisors 1 and n. Hence $\tau(n)$ can never be less than 2 for $n > 1$.
II. Let "p" be a prime. We know that the prime p has only two divisors 1 and p itself. Hence $\tau(p) = 2$.

It follows that $\tau(n)$ assumes the value 2 for infinitely many values of n because the number of prime is infinite.

Thus we write

$$\lim_{n\to\infty} \tau(n) = 2$$

III. If $n= p^a$, p is a prime, Then $\tau(n) = a + 1$

Therefore $\tau(n)$ assumes values larger than any given number for sufficiently large values of a.

Hence we may write

$$\lim_{n\to\infty} \tau(n) = \infty$$

The function $\tau(n)$ thus behaves in a very irregular way.

Behaviour of σ(n) as he n $\to \infty$

We know that $\sigma(n) > n$ for all $n > 1$. It follows $\sigma(n)$ assumes values larger than any given number for sufficiently large values of n. Thus we have

$$\lim_{n\to\infty} \tau(n)$$

$$= \infty$$

On the other hand $\sigma(n)$ is not a monotonic increasing functions of n.

2.20 Theorem. Let p be a prime > 3. Then $\sigma(p-1) > \sigma(p)$.

SJIF Impact Factor (2025): 8.688| ISI I.F. Value: 1.241| Journal DOI: 10.36713/epra2016 ISSN: 2455-7838(Online)
EPRA International Journal of Research and Development (IJRD)
Volume: 10 | Issue: 5 | May 2025 - Peer Reviewed Journal

Proof.

I. Let $p-1=2^t, t>1. Then$
$\sigma(\mathrm{p}-1)=\sigma(2^t)=2^t+2^{t-1}+\cdots+2+1$

Hence

$\sigma(p-1)>\ 2^t+2>p+1=\ \sigma(p).$

II. Let $(p-1)=p1^{a1}p2^{a2}\ldots\ldots\ pk^{ak}.$ $Then$

$\sigma(p-1)=\sigma(p1^{a1})\,\sigma(p2^{a2})\ldots\ldots\ldots\ldots\ldots\sigma(pk^{ak})$

$\geq(p1^{a1}+1)(p2^{a2}+1)\ldots\ldots\ldots\ldots(p1^{a1}+1)$

$\geq p1^{a1}p2^{a2}\ldots\ldots\ldots\ldots pk^{ak}+(p1^{a1}+p2^{a2}\ldots\ldots+pk^{ak})$

$\geq p-1+k+1$

$>p+1$ $Since\ k>1$

$=\sigma(p)$

Thus $\sigma(n)$ is an irregular function of n although not as $\tau(n)$

REFERENCES

1. *S. G. Telang, Number theory, Tata Mc grow –Hill Publishing Company Limited New Delhi, 1996.*
2. *David M. Burton, Elementary Number Thoery, Tata Mc grow-Hill publishing company limited, New Delhi, 2006.*
3. *Thomas Koshy, Elementary Number Thoery with Applications, Academic Press is an imprint of Elsevier.*